

Cardiac arrest after severe traumatic brain injury can be survivable with good outcomes

Zirun Zhao , Justine J Liang, Zhe Wang, Nathan J Winans, Matthew Morris, Stephen Doyle, Adam Fry, Susan M Fiore, Sima Mofakham, Charles B Mikell

Department of Neurosurgery,
Stony Brook University
Renaissance School of Medicine,
Stony Brook, New York, USA

Correspondence to

Zirun Zhao; Zirun.zhao@stonybrookmedicine.edu

Received 4 November 2020
Revised 12 December 2020
Accepted 17 January 2021

ABSTRACT

Background Resuscitation for traumatic cardiac arrest (TCA) in patients with severe traumatic brain injury (sTBI) has historically been considered futile. There is little information on the characteristics and outcomes of these patients to guide intervention and prognosis. The purpose of the current study is to report the clinical characteristics, survival, and long-term neurological outcomes in patients who experienced TCA after sTBI and analyze the factors contributing to survival.

Methods A retrospective review identified 42 patients with TCA from a total of 402 patients with sTBI (Glasgow Coma Scale (GCS) score ≤ 8) who were admitted to Stony Brook University Hospital, a level I trauma center, from January 2011 to December 2018. Patient demographics, clinical characteristics, survival, and neurological functioning during hospitalization and at follow-up visits were collected.

Results Of the 42 patients, the average age was 45 years and 21.4% were female. Eight patients survived the injury (19.0%) to discharge and seven survived with good neurological function. Admission GCS score and bilateral pupil reactivity were found to be significant indicators of survival. The mean GCS score was 5.3 in survivors and 3.2 in non-survivors ($p=0.020$). Age, Injury Severity Score, or cardiac rhythm was not associated with survival. Frequent neuroimaging findings included subarachnoid hemorrhage, subdural hematoma, and diffuse axonal injury.

Discussion TCA after sTBI is survivable and seven out of eight patients in our study recovered with good neurological function. GCS score and pupil reactivity are the best indicators of survival. Our results suggest that due to the possibility of recovery, resuscitation and neurosurgical care should not be withheld from this patient population.

Level of evidence Level IV, therapeutic/care management.

INTRODUCTION

Traumatic brain injury (TBI) is a major cause of morbidity and mortality worldwide.¹ Severe TBI (sTBI), defined as Glasgow Coma Scale (GCS) score equal or less than 8,² often occurs in the setting of profound overall injury. One feared consequence of major trauma is traumatic cardiac arrest (TCA), which has a survival rate of less than 10% despite resuscitation by trained medical professionals.³ Because of the extremely poor prognosis, resuscitation of patients with TCA has been historically considered futile.³⁻⁵

The effect of cardiac arrest on the brain is well characterized. Much of the brain injury is not due to the initial insult, but rather to the inflammatory cascade that follows the initial cessation of cerebral blood flow.⁶ Persistent neurological deficits often result from secondary brain injury and neuronal cell death, which occur after the initial anoxic period.⁷ As a result, hypoxic-ischemic brain injury after cardiac arrest often results in long-term disability in survivors. However, reports of cardiac arrest after head trauma are extremely rare. Although studies of TCA from all trauma included some cases of TBI, the extent of their brain injury, neuroimaging characteristics, and more importantly long-term neurological function remain unclear.

Given complications that arise from cardiac arrest in addition to TBI, it is challenging to recommend neurosurgical interventions and provide prognostic information to families. Currently, it is thought that there is little chance of adequate recovery,³⁻⁵ but our study suggests that this may not necessarily be the case. The current study aims to characterize the clinical features, long-term survival and neurological outcomes, and prognostic factors for survival in patients who experienced cardiac arrest after sTBI. Our study would ultimately serve to provide information on management of patients with sTBI with cardiac arrest, who have been historically managed with extremely poor prognosis.

PATIENTS AND METHODS

Study design and data collection

We conducted a retrospective chart review at Stony Brook University Hospital, a level I trauma Center in Long Island, New York. Patient consent waivers were obtained from the Institutional Review Board office. We included all 402 adult patients (aged ≥ 18 years) during a period of 8 years (January 2011 and December 2018) who sustained sTBI, defined by a GCS score of 8 or less on presentation to the emergency department (ED). We then identified 42 patients who experienced TCA either outside of the hospital or at the ED. We excluded patients (1) with penetrating trauma, (2) who did not have TCA, (3) whose cardiac arrest happened after hospital admission or secondary to medical reasons (one patient had cardiac arrest during dialysis and subsequently fell), and (4) with significant cardiac comorbidities. We also excluded 11 patients who met the above criteria but whose care was withdrawn due to extremely poor functional outcome. All data were extracted from the hospital electronic medical record (Cerner Millennium, PowerChart).

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Zhao Z, Liang JJ, Wang Z, et al. *Trauma Surg Acute Care Open* 2021;**6**:e000638.

Data were collected on age, sex, mechanism of injury, Injury Severity Score (ISS), GCS score on ED presentation, preintervention non-contrast head CT scans, initial pupil diameter and reactivity to light, neurosurgical interventions (external ventricular drain placement, hematoma evacuation, decompressive craniectomy), and length of stay. Regarding TCA, location (on the scene, transit, or at the ED), arrest rhythm, return of spontaneous circulation (ROSC), and airway management outside of the hospital (endotracheal intubation, bag mask ventilation, or laryngeal mask airway) were also recorded.

ISS was computed by summing the square of the three highest individual Abbreviated Injury Scale (AIS) scores. Head AIS score was listed separately. Non-contrast CT scan at admission was graded by the 6-point Rotterdam Scoring System.⁸ The Rotterdam scores evaluate injury severity based on the presence of basal cistern effacement, midline shift of >5 mm, epidural mass, and intraventricular or subarachnoid hemorrhage.

Outcomes

The primary outcome of the study is the survival status after TCA following sTBI. The surviving patients (n=8) were further evaluated at 6-month and 1-year postdischarge follow-up in the outpatient clinic. Data were obtained from clinical notes at these follow-up visits. Cases were scored according to Glasgow Outcome Scale-Extended and a modified Rankin Scale (mRS) for neurological disability at their follow-up appointments from 6 months to 1 year after discharge.⁹

Statistical analysis

Categorical variables were described as frequencies and percentages and were compared using χ^2 test or Fisher's exact test. Numerical variables were presented as mean with SD and compared using Student's t-test. Variables with p values less than 0.05 were considered significant. All analyses were performed in SPSS V.26.

RESULTS

Patient characteristics

Of the 402 adult cases of sTBI at Stony Brook University Hospital from 2011 to 2018, 42 patients of TCA were included in the study, with 8 surviving patients at discharge (figure 1). As shown in table 1, of the 42 patients, the age range was 18 to 84 years, with a mean of 45 years. Of these patients 21.4% were female. The patients in this study suffered from severe head




Figure 1 Flow chart of patient selection. 42 patients with traumatic cardiac arrest (TCA) in the current study were selected from 402 patients with severe traumatic brain injury (sTBI) admitted to Stony Brook University Hospital from January 2011 to December 2018. Exclusion criteria were listed. The mean Glasgow Coma Scale (GCS) scores of the surviving and deceased patients with TCA were 5.3 and 3.2, respectively.

Table 1 Demographics and clinical characteristics of reviewed patients

Demographics	
Age, mean±SD	44.5±20.2
Sex	
Female (%)	9 (21.4)
Male (%)	33 (78.6)
ISS, mean±SD	38.2±14.0
Head AIS score, mean±SD	4.2±1.1
GCS score, mean±SD	3.5±1.3
GCS motor score, mean±SD	1.4±0.9
Mechanism of injury (%)	
Fall	6 (14.3)
Motor vehicle collision	12 (28.6)
Pedestrian struck	15 (35.7)
Motorcycle crash	9 (21.4)
Airway management (%)	
Endotracheal intubation	30 (71.4)
Bag mask ventilation	2 (4.8)
Cardiac arrest location (%)	
On the field	27 (64.3)
En route	10 (23.8)
ED	5 (11.9)
Arrest rhythm (%)	
Asystole	16 (38.1)
PEA	15 (35.7)
Ventricular arrhythmia	2 (4.8)
Unknown	9 (21.4)
Imaging findings (%)	
Subdural hematoma	11 (42.3)
SAH	21 (80.7)
Hypoxic injury	8 (30.8)
Herniation	6 (23.1)
Rotterdam score, mean	2.8±1.2
Neurosurgical interventions (%)	
Bolt	11 (26.2)
EVD	5 (11.9)
Decompressive craniectomy	2 (4.8)
Cause of death (%)	
Cardiac arrest	27 (64.3)
Brain death	7 (16.7)

AIS, Abbreviated Injury Scale; ED, emergency department; EVD, external ventricular drain; GCS, Glasgow Coma Scale; ISS, Injury Severity Score; PEA, pulseless electrical activity; SAH, subarachnoid hemorrhage.

trauma with a mean GCS score of 3.5 and severe overall injury with a mean ISS of 38.2. The leading mechanism of injury was pedestrian struck (35.7%), followed by motor vehicle collision (28.6%), motorcycle crash (21.4%), and fall (14.3%).

All 42 patients experienced cardiac arrest related to trauma, either prior to arriving at the hospital or very shortly after presentation to the ED. Figure 2 illustrates the cardiac rhythms before cardiopulmonary resuscitation (CPR). The most common rhythm was asystole (38.1%), followed by pulseless electrical activity (PEA) (35.7%) and ventricular arrhythmia (4.8%). Regarding emergent airway management, 30 (71.4%) patients




Figure 2 Cardiac rhythms before initiation of cardiopulmonary resuscitation. The most common cardiac rhythm is asystole (38.1%), followed by pulseless electrical activity (PEA) (35.7%).

were intubated at the scene of injury and two (4.8%) received bag mask ventilation.

The survival rate in patients who experienced TCA after sTBI was 19.0% (8 out of 42). In the 34 patients who did not survive, the most common cause of death was cardiac arrest in 27 patients (64.3%), followed by brain death in 7 (16.7%).

Characteristics of survivors

Of the total 42 patients with TCA in this study, eight survived to discharge (table 2). The mean age of the survivors was 38.6 years (range, 18–66), with two women and six men. The average length of stay was 53.3 days (range, 9–100). Three (37.5%) of these patients were pedestrians struck at high speeds, three

(37.5%) were in motor vehicle crash, and two were motorcyclists struck by cars (25%). These cases all had additional injuries apart from sTBI, demonstrated by a mean ISS of 33.6. Two patients achieved ROSC under 5 minutes. All eight patients had no known cardiac comorbidities prior to sTBI presentation. All survivors followed commands during their hospitalization. The average time to command following was 16.1 days (range 2–36 days).

Seven out of eight patients survived with good neurological function, as demonstrated by an mRS score of equal or less than 4 at their follow-up visits. At 6-month to 1-year follow-up, two patients had good recovery with an mRS score of 1. They were able to carry out activities of daily living independently, although with residual symptoms (right eye vision loss in patient 5). Two patients had slight disability but were able to carry out daily activities with minimal assistance. Three patients had moderate to severe disability (paraplegia) but had the ability to communicate and/or follow commands. One patient scored 5 on the mRS, requiring constant nursing care, as well as a tracheotomy and a gastrostomy.

Factors contributing to survival

We compared the clinical and imaging characteristics of survivors and non-survivors (table 3). Survivors had higher overall GCS score ($p=0.020$) and its motor component ($p=0.026$) compared with non-survivors. Pupil reactivity on ED presentation was significantly different between survivors and non-survivors ($p=0.001$) (figure 3). The majority of non-survivors had non-reactive and dilated pupils, although three of them had bilateral reactivity on initial presentation. This was in contrast

Table 2 Clinical characteristics of survivors

Patient number	Age (range)	MOI	Admission GCS score (motor), ISS	Time to ROSC, type of arrest	Pupil reactivity	Brain injuries	Follow-up (6 months–1 year)	Modified Rankin score, GOS-E score
1	20s	Motorcycle crash	6, 45	Unknown, PEA	Bilateral reactivity	Multicompartmental intracranial hemorrhage. Epidural hematoma. Effacement of suprasellar cisterns.	Able to walk. Neurologically intact.	2, 8
2	20s	Motor vehicle collision	3, 22	Unknown, unknown	Non-reactive, constricted	Left SAH with bifrontal contusions. Cerebral edema.	Able to walk. Able to carry out ADLs.	2, 7
3	10s	Pedestrian struck	3, 33	Unknown, PEA	Bilateral reactivity	Diffuse SAH. Cerebral edema. Sulcal effacement. Basilar cisterns effacement.	Requires total assistance for ADLs. Dependent on tracheotomy and gastric tube.	5, 3
4	60s	Pedestrian struck	8, 26	4 min, PEA	Bilateral reactivity	Intracranial hemorrhage. SAH.	Able to carry out ADLs independently.	1, 6
5	50s	Motorcycle crash	6, 42	2 min, PEA	Bilateral reactivity	Bilateral SAH.	Able to walk with prosthesis. Able to communicate. Residual right eye vision loss.	1, 5
6	50s	Motor vehicle collision	8, 50	Unknown, asystole	Bilateral reactivity	SAH in posterior right frontal regions. Contusions in anterior frontal lobe.	Paraplegia but otherwise able to communicate.	4, 4
7	50s	Motor vehicle collision	5, 17	Unknown, unknown	Non-reactive, constricted	Diffuse SAH bilaterally. Intraventricular hemorrhage.	Non-verbal but followed commands. Quadriplegic. Dependent on tracheotomy and gastric tube.	4, 3
8	20s	Pedestrian struck	3, 34	Unknown, PEA	Bilateral reactivity	Right-sided pneumocephalus. Left frontal lobe hemorrhagic contusion. Bilateral frontal SAH.	Able to communicate. Regained use of arms. Dependent on tracheotomy and gastric tube.	4, 4

ADLs, activities of daily living; GCS, Glasgow Coma Scale; GOS-E, Glasgow Outcome Scale-Extended; ISS, Injury Severity Score; MOI, mechanism of injury; PEA, pulseless electric activity; ROSC, return to spontaneous circulation; SAH, subarachnoid hemorrhage.

Table 3 Comparisons between survivors and non-survivors of traumatic arrest after TBI

	Survivors (n=8)	Non-survivors (n=34)	P value
Age, mean±SD	38.6±18.7	45.8±20.8	0.286
Sex			0.784
Female (%)	2 (25)	7 (20.6)	
Male (%)	6 (75)	27 (79.4)	
ISS, mean±SD	33.6±11.6	39.3±14.4	0.306
Head AIS score, mean±SD	3.9±1.1	4.3±1.1	0.365
GCS score, mean±SD	5.3±2.1	3.2±0.5	0.020
GCS motor score, mean±SD	2.5±1.3	1.1±0.4	0.026
Pupil diameter and reactivity (%)			0.001
Bilateral reactivity	5 (62.5)	3 (8.8)	
Bilateral constriction and non-reactivity	2 (25)	3 (8.8)	
Bilateral dilation and non-reactivity	1 (12.5)	27 (79.4)	
Unilateral dilation and non-reactivity	0	1 (2.9)	
Positive FAST examination (%)	2 (66.7)	19 (70.4)	0.894
Airway management (%)			0.139
Intubation	7 (87.5)	23 (67.6)	
Bag ventilation	1 (12.5)	1 (2.9)	
Cardiac arrest location (%)			0.701
On the field	6 (75)	21 (61.8)	
En route	1 (12.5)	9 (26.5)	
ED	1 (12.5)	4 (11.8)	
Arrest rhythm (%)			0.378
Asystole	1 (12.5)	15 (44.1)	
PEA	5 (62.5)	10 (29.4)	
Ventricular arrhythmia	0	2 (5.8)	
Unknown	2 (25)	7 (20.6)	
Mechanism of injury (%)			0.625
Fall	0	6 (17.6)	
MVC	3 (37.5)	9 (26.5)	
Pedestrian vs auto	3 (37.5)	12 (35.3)	
Motorcycle crash	2 (25)	7 (20.6)	
Violence	0	1 (2.2)	
Imaging findings (%)			
Subdural hematoma	1 (12.5)	10 (55.6)	0.001
SAH	8 (100)	13 (72.2)	0.007
Hypoxic injury	2 (25)	6 (33.33)	0.029
Herniation	1 (12.5)	5 (27.8)	0.019
Rotterdam score, mean±SD	2.63±0.9	2.89±1.4	0.625
Interventions (%)			
ICP monitoring	7 (87.5)	4 (11.8)	<0.001
EVD	3 (37.5)	2 (5.9)	0.060
Decompressive craniectomy	0	2 (5.9)	0.594

Bolded: p<0.05

AIS, Abbreviated Injury Scale; ED, emergency department; EVD, external ventricular drain; FAST, focused assessment with sonography for trauma; GCS, Glasgow Coma Scale; ICP, intracranial pressure; ISS, Injury Severity Score; MVC, motor vehicle collision; PEA, pulseless electric activity; SAH, subarachnoid hemorrhage; TBI, traumatic brain injury.

to survivors, of whom the majority had bilateral pupil reactivity, with only two who presented with non-reactive and constricted pupils. Patients who presented with bilateral reactivity at admission had a survival rate of 62.5%, whereas those presenting with non-reactive pupils had a survival rate of only 8.8%. There was




Figure 3 Pupil diameter and reactivity in patients with traumatic brain injury with traumatic cardiac arrest are compared between survivors and non-survivors in four categories.

no significant difference in age, gender, ISS, or mechanisms of injury between survivors and non-survivors. Similarly, the location of CPR initiation, intubation status prior to hospital arrival, or traumatic arrest rhythm did not contribute significantly to mortality.

Neuroimaging and neurosurgical interventions

Non-contrast head CT was performed in 26 out of 42 patients with TCA. The rest died soon after ED arrival, before they could be scanned. Of the patients 21 (80.7%) showed evidence of subarachnoid hemorrhage, 11 (42.3%) had subdural hematoma, and 8 (30.8%) showed diffuse hypoxic injury with poor gray-white matter differentiation. Seven patients had evidence of herniation on neuroimaging, including transtentorial, subfalcine, and tonsillar herniation. The mean Rotterdam score was 2.8.

Interestingly, 87.5% of the survivors received intracranial pressure monitoring, significantly more than 11.8% in the non-survivors (p<0.001). This was likely due to patients who did not survive dying rapidly from the injuries, neurosurgery interventions being considered futile, or risks outweighing the benefits. Of the patients 16.7% underwent neurosurgical procedures. Seven patients received external ventricular drainage and two underwent decompressive craniectomy.

DISCUSSION

Resuscitation of patients with TCA has a low success rate and is traditionally considered futile.³⁻⁵ Little is known about patients with TCA after sTBI, which is a major cause of death and disability in itself. Our study demonstrates that, despite the grim odds of survival and extensive injury, some patients do survive, even with good neurological function. Although limited by the small sample size, the current study has a survival rate of 19.0%, which is comparable with cardiac arrest associated with all other trauma, and even as good or better than outcomes from out-of-hospital cardiac arrest of any cause.^{10 11} It is important to note that some of our cases of survival achieved ROSC less than 5 minutes, which may contribute to the positive neurological outcomes observed. Our results show that TBI may not substantially affect TCA survival, and timely resuscitation can yield good neurological outcomes.

The two most important clinical features that provide prognostic information in our study are GCS score and pupil reactivity. GCS is routinely used to evaluate neurological function in patients who have sustained head injuries. We showed that GCS score is a significant predictor of survival after cardiac arrest associated with sTBI. GCS motor score is also significantly

higher in survivors compared with non-survivors. This result is consistent with previous studies,^{12,13} including the CRASH trial,¹⁴ which demonstrated strong correlation between admission GCS score with mortality and morbidity at discharge and at 6-month follow-up. The GCS score in this study is recorded on arrival at the ED, which has been shown to have higher predictive value than those obtained on the scene as a result of initial management of airway, breathing, and circulation.¹³ On the contrary, ISS was not a significant predictor in our patient population. This is partly because all of our patients have very severe overall injury (ISS >15). The results of our study suggest that neurological function assessment by GCS outweighs overall injury in providing prognostic information in sTBI.

Pupil diameter and reactivity to light provide vital information on the extent of brain damage. A dilated unreactive pupil indicates compression of the oculomotor nerve, often due to midline shift and uncal herniation, and thus worse prognosis. This was reflected in our results that most of the patients who survived had bilaterally reactive pupils, whereas almost all of those with non-reactive and dilated pupils did not survive. Previous report has also shown that pupillary reactivity is a reliable prognostic indicator of mortality in sTBI both on the field and at admission, compared with GCS, making it a more useful early indicator of survival.¹⁵ One limitation of pupil reactivity is that it could be affected by sedative medications. However, most of our patients only had light sedation on arrival to the ED. It is important to note that previous studies have made the effort to combine GCS score and pupil response to provide more accurate prognosis of patients with sTBI, highlighting the importance of both clinical assessments in evaluating severe brain trauma.⁸ Our study is in concordance with these findings in that GCS score and pupil response are the single two most important indicators of survival.

The most common initial rhythm before cardiac arrest in our study is PEA, followed by asystole. These two cardiac rhythms are most commonly associated with obstructive, hypovolemic, and cardiogenic shock as well as tissue hypoxia.^{16,17} Although closely related to traumatic injury, no specific cardiac rhythm was significantly associated with mortality in our study. Other general clinical characteristics related to trauma, such as mechanisms of injury, intubation status on the scene, and the location of cardiac arrest, also did not significantly contribute to mortality in our cohort, which is consistent with current literature on TCA with larger sample sizes.^{10,16,18} Studies focusing on the relationship between age and mortality from TCA have mixed findings. Majority shows that older age is associated with worse outcome.^{11,19} In our study, however, there was no difference in age between patients who survived and those who did not. This result could be attributed to the fact that we excluded patients with severe chronic comorbidities and cardiac arrest due to medical causes. Both of these conditions have increased incidence with age and forebode worse outcomes seen in elderly patients.^{20,21}

Several consequences of brain trauma after TCA are evidenced on imaging. In our study, most patients who underwent CT scans had radiological evidence of subarachnoid hemorrhage and subdural hematoma. Other frequent neuroimaging findings in patients include diffuse axonal injury with poor gray-white matter differentiation. Hypoxic-ischemic brain injury is one of the most serious consequences of cardiac arrest, reported to be present in one-fifth of all out-of-hospital cardiac arrest cases.^{6,22} The effects of cardiac arrest on brain tissue are twofold—a primary injury immediately as a result of cessation of cerebral blood flow and a secondary injury related to reperfusion, which takes hours or days to occur. Whether the brain imaging findings in our study

are attributed to the primary or the secondary mechanisms is not clear. Previous mechanistic studies have found common biomarkers of cardiac arrest and TBI, which may suggest shared pathophysiological cascades in these two insults.²³ Future studies are needed to elucidate the biochemical changes in the brain tissue in cases of TCA after sTBI.

Our study is a retrospective chart review and is subject to limitations of the study design. The sample size is small, partially limited by the rare event of cardiac arrest after sTBI. We recognize that GCS score at admission may be affected by confounding effects from sedation, neuromuscular blockade, and rapid sequence intubation during resuscitation on the field, as well as inter-rater discrepancy. Nevertheless, GCS remains an essential clinical tool for evaluating the level of consciousness when patients with severe trauma arrive at the hospital. We did not include the underlying health status of the patients in the analysis, which could be a factor contributing to their survival and recovery. However, we did exclude those with significant cardiac comorbidity and those with cardiac arrest due to other medical reasons. Limited by the nature of chart review study, we did not analyze the transport time from cardiac arrest to ED arrival, CPR duration, or time to definitive airway, which could also impact the success of resuscitation. Nonetheless, we think that the reported data are representative of the TCA population in general.

In conclusion, this study demonstrates that patients with cardiac arrest after sTBI do survive and can recover with decent neurological function, although the mortality rate remains high. Initial clinical features including GCS score, pupillary diameter, and reactivity to light were important indicators of survival. Some frequent neuroimaging findings on non-contrast head CT in this unique group of patients include subarachnoid hemorrhage, subdural hematoma, and hypoxic axonal injury. Given the chance of survival and neurological recovery, cardiac arrest event should not exclude patients with sTBI from getting proper resuscitation and appropriate neurosurgical care.

Acknowledgements We would like to thank Jane McCormack, Emily Huang, and the Stony Brook Trauma, Emergency Surgery, and Surgical Critical Care Division faculty and staff for their assistance in this study.

Contributors Conception and design: CBM, ZZ, JLL, SM. Acquisition of data: CBM, ZZ, JLL, NJW, MM, SD, AF. Analysis and interpretation of data: CBM, ZZ, JLL, ZW, NJW, MM, SD, AF, SMF, SM. Drafting the article: CBM, ZZ, JLL, SM. Critically revising the article: CBM, ZZ, JLL, ZW, SM. Approved the final version of the article on behalf of all authors: CBM. Statistical analysis: CBM, ZZ, JLL. Administrative/technical/material support: CBM, ZZ, JLL. Study supervision: CBM, SM.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study was approved by the Stony Brook Institutional Review Board (IRB-2019-00464).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The data set used in this manuscript is part of a larger traumatic brain injury database at our institution. We currently do not have the approval from IRB to share any portion of the data set as it is protected patient information. Please contact the corresponding author for data set requests at this time.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Zirun Zhao <http://orcid.org/0000-0001-9411-0224>

REFERENCES

- 1 Stocchetti N, Carbonara M, Citerio G, Ercole A, Skrifvars MB, Smielewski P, Zoerle T, Menon DK. Severe traumatic brain injury: targeted management in the intensive care unit. *Lancet Neurol* 2017;16:452–64.
- 2 Picetti E, Iaccarino C, Servadei F. Letter: guidelines for the management of severe traumatic brain injury fourth edition. *Neurosurgery* 2017;81:E2–E.
- 3 Rosemurgy AS, Norris PA, Olson SM, Hurst JM, Albrink MH. Prehospital traumatic cardiac arrest: the cost of futility. *J Trauma* 1993;35:468–74.
- 4 Millin MG, Galvagno SM, Khandker SR, Malki A, Bulger EM. Withholding and termination of resuscitation of adult cardiopulmonary arrest secondary to trauma: resource document to the joint NAEMSP-ACSCOT position statements. *J Trauma Acute Care Surg* 2013;75:459–67.
- 5 Battistella F *et al.* Field triage of the pulseless trauma patient. *Arch Surg* 1999;134:742–6.
- 6 Sekhon MS, Ainslie PN, Griesdale DE. Clinical pathophysiology of hypoxic ischemic brain injury after cardiac arrest: a “two-hit” model. *Critical Care* 2017;21:1–10.
- 7 Meaney DF, Morrison B, Dale Bass C. The mechanics of traumatic brain injury: a review of what we know and what we need to know for reducing its societal burden. *J Biomech Eng* 2014;136.
- 8 Steyerberg EW, Mushkudiani N, Perel P, Butcher I, Lu J, McHugh GS, Murray GD, Marmarou A, Roberts I, Habbema JDF, *et al.* Predicting outcome after traumatic brain injury: development and international validation of prognostic scores based on admission characteristics. *PLoS Med* 2008;5:e165.
- 9 Quinn TJ, Dawson J, Walters MR, Lees KR. Reliability of the modified Rankin scale: a systematic review. *Stroke* 2009;40:3393–5.
- 10 Lockey D, Crewdson K, Davies G. Traumatic cardiac arrest: who are the survivors? *Ann Emerg Med* 2006;48:240–4.
- 11 Stiell IG, Wells GA, Field B, Spaite DW, Nesbitt LP, De Maio VJ, Nichol G, Cousineau D, Blackburn J, Munkley D, *et al.* Advanced cardiac life support in out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:647–56.
- 12 Kung W-M, Tsai S-H, Chiu W-T, Hung K-S, Wang S-P, Lin J-W, Lin M-S. Correlation between Glasgow coma score components and survival in patients with traumatic brain injury. *Injury* 2011;42:940–4.
- 13 Lesko MM, Jenks T, O'Brien SJ, Childs C, Bouamra O, Woodford M, Lecky F. Comparing model performance for survival prediction using total Glasgow coma scale and its components in traumatic brain injury. *J Neurotrauma* 2013;30:17–22.
- 14 . MRC CRASH Trial Collaborators. Predicting outcome after traumatic brain injury: practical prognostic models based on large cohort of international patients. *BMJ* 2008;336:425–9.
- 15 Majdan M, Steyerberg EW, Nieboer D, Mauritz W, Rusnak M, Lingsma HF. Glasgow coma scale motor score and pupillary reaction to predict six-month mortality in patients with traumatic brain injury: comparison of field and admission assessment. *J Neurotrauma* 2015;32:101–8.
- 16 Konesky KL, Guo WA. Revisiting traumatic cardiac arrest: should CPR be initiated? *Eur J Trauma Emerg Surg* 2018;44:903–8.
- 17 Morrison JJ, Poon H, Rasmussen TE, Khan MA, Midwinter MJ, Blackburne LH, Garner JP. Resuscitative thoracotomy following wartime injury. *J Trauma Acute Care Surg* 2013;74:825–9.
- 18 Evans CCD, Petersen A, Meier EN, Buick JE, Schreiber M, Kannas D, Austin MA. Prehospital traumatic cardiac arrest: management and outcomes from the resuscitation outcomes Consortium epistry-trauma and PROPHET registries. *J Trauma Acute Care Surg* 2016;81:285.
- 19 Baker SP, O'Neill B, Haddon W, Long WB. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma* 1974;14:187–96.
- 20 Ferraris VA, Ferraris SP, Saha SP. The relationship between mortality and preexisting cardiac disease in 5,971 trauma patients. *J Trauma* 2010;69:645–52.
- 21 McMahon DJ, Schwab CW, Kauder D. Comorbidity and the elderly trauma patient. *World J Surg* 1996;20:1113–20.
- 22 Laver S, Farrow C, Turner D, Nolan J. Mode of death after admission to an intensive care unit following cardiac arrest. *Intensive Care Med* 2004;30:2126–8.
- 23 Mussack T, Biberthaler P, Kanz K-G, Wiedemann E, Gippner-Steppert C, Mutschler W, Jochum M. Serum S-100b and interleukin-8 as predictive markers for comparative neurologic outcome analysis of patients after cardiac arrest and severe traumatic brain injury. *Crit Care Med* 2002;30:2669–74.