

Percutaneous thoracostomy for traumatic hemothoraces: a call for research

Marc de Moya 

To cite: de Moya M. Percutaneous thoracostomy for traumatic hemothoraces: a call for research. *Trauma Surg Acute Care Open* 2024;**9**:e001384. doi:10.1136/tsaco-2024-001384



► <http://dx.doi.org/10.1136/tsaco-2023-001298>

© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Surgery, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

Correspondence to

Dr Marc de Moya; mdemoya@mcw.edu

McLauchlan and others report a case series of nine trauma patients with hemothoraces that were drained with a percutaneous 14Fr drain and lavaged.¹ They demonstrate the feasibility of performing thoracic lavage via the 14Fr drain. In doing so, they were able to, in theory, leverage the benefits of a smaller percutaneous tube (less pain)² with the benefits of lavage (lower incidence of retained hemothoraces).³ It is the first report of combining the techniques and shows promise. There are a number of questions that it raises which certainly will need to be evaluated with a larger prospective cohort. The type of percutaneous tube used was one of the pigtail variety which may need to be evaluated in comparison to its straight tube variety. The timing of the lavage and the possible need for a repeat lavage will need to be addressed. There may be a role for lavage with a pigtail that is image guided for empyema or the installation of lytic therapy. Of note, the authors were able to quantify the hemothoraces in some of the patients using the Mergo formula calculated on a CT scan prior to tube placement. The utilization of a more objective measure of the volume of the hemothorax is another evolving field that warrants a closer examination.

Overall, the authors offer a feasible technique for lavaging the chest with less pain and may be a promising approach to the stable patient with a traumatic hemothorax. I look forward to future studies that explore the many questions regarding the management of hemothoraces that remain.

Contributors MD is the sole author.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Provenance and peer review Commissioned; internally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Marc de Moya <http://orcid.org/0000-0002-1942-4650>

REFERENCES

- 1 Ali N, Beyer C, Brinson M, Joergensen S, Yelon J, Dumas R, Vella M, Cannon J. Percutaneous Thoracostomy with Thoracic Lavage for traumatic Hemothorax: a performance improvement initiative. *TSACO* 2024.
- 2 Kulvatunyou N, Erickson L, Vijayasekaran A, Gries L, Joseph B, Friese RF, O'Keeffe T, Tang AL, Wynne JL, Rhee P. Randomized clinical trial of pigtail catheter versus chest tube in injured patients with uncomplicated traumatic pneumothorax. *Br J Surg* 2014;101:17–22.
- 3 Kugler NW, Milia D, Paul JS, TW. Thoracic irrigation prevents retained hemothorax: a prospective propensity scored analysis. *J Trauma Acute Care Surg* 2017;83:1136–41.