Different strokes: differences in the characteristics and outcomes of BCVI and non-BCVI strokes in trauma patients

Michelle K McNutt,¹ Cedar Slovacek,² David Rosenbaum,³ Hari Kishan Reddy Indupuru,² Xu Zhang,⁴ Bryan A Cotton,¹ John Harvin ,¹ Charles E Wade,¹ Sean I Savitz,⁵ Lillian S Kao¹

¹Surgery, University of Texas McGovern Medical School, Houston, Texas, USA

²University of Texas McGovern Medical School, Houston, Texas, USA

³Neurosurgery, University of Texas McGovern Medical School, Houston, Texas, USA

⁴Center for Clinical and Translational Sciences, University of Texas McGovern Medical School, Houston, Texas, USA

⁵UTHealth Institute for Stroke and Cerebrovascular Disease, University of Texas McGovern Medical School, Houston, Texas, USA

Correspondence to

Dr Michelle K McNutt, Surgery, University of Texas Health Science Center at Houston, Houston, TX 77030, USA; Michelle.K.McNutt@uth.tmc.edu

Received 4 March 2020

Revised 14 April 2020

Accepted 21 April 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: McNutt MK, Slovacek C, Rosenbaum D, et al. *Trauma Surg Acute Care Open* 2020;**5**:e000457.

ABSTRACT

Background Although strokes are rare in trauma patients, they are associated with worse functional and cognitive outcomes and decreased mobility. Blunt cerebrovascular injury (BCVI)-related strokes and mortality have decreased, likely due to refined screening and treatment algorithms in trauma literature; however, there is a paucity of research addressing non-BCVI strokes in trauma. The purpose of this study is to evaluate the incidence, etiology, and risk factors of stroke in our trauma population in order to identify preventive strategies.

Methods This study was a retrospective review of all adult trauma patients admitted to a level 1 trauma hospital who suffered a stroke during trauma admission from 2010 to 2017. Data were collected from the prospectively maintained trauma and stroke databases. Stroke etiology was determined by a vascular neurologist.

Results Of the 43 674 adult trauma patients admitted during the study period, 99 (0.2%) were diagnosed with a stroke during the index admission. Twenty-one (21%) strokes were due to BCVI. Seventy-eight (79%) strokes were due to non-BCVI etiologies. Patients with non-BCVI strokes were older, less severely injured, and had more medical comorbidities compared with patients with a BCVI stroke. While patients with a BCVI stroke were more likely to suffer multiple traumatic injuries from MVC (76% vs 28%, $p<0.001$), non-BCVI strokes had more isolated extremity injuries from fall mechanism (55% vs 10%, $p<0.001$). Over the study period, the age and incidence of stroke and BCVI ($p<0.001$) increased. However, the rate of BCVI strokes decreased while the rate of non-BCVI strokes increased.

Discussion The incidence of stroke has increased despite aggressive screening and treatment of BCVI. This increase is primarily due to non-BCVI strokes which are associated with advanced age and medical comorbidities after low mechanism traumatic injury. Medical optimization of comorbid conditions during trauma hospitalization will become increasingly important for stroke prevention as the population ages.

Level of evidence: Level III

BACKGROUND

Blunt cerebrovascular injuries (BCVIs) after trauma have traditionally been a rare diagnosis, occurring in 1%–2% of all blunt trauma patients. However, with expanded screening criteria and improved

diagnostic modalities,^{1–11} the incidence of BCVI has increased over the past three decades.¹² Despite this increase, BCVI-related strokes and mortality have decreased,¹² likely as a result of both improved diagnostic and treatment algorithms.^{12–15}

Although strokes are rare after trauma, they can have significant short and long-term impact on functional and cognitive status. In a study from the Netherlands, Hundersmarck *et al* demonstrated that patients with BCVIs have worse long-term functional outcomes than those in the general trauma population.¹⁶ In addition, patients with BCVI and ischemic stroke had worse long-term outcomes in terms of mobility and self-care than patients with BCVI alone. Patients with BCVI and traumatic brain injury had worse cognitive outcomes. Shahan *et al* demonstrated that stroke, along with age, predicts decreased mobility after BCVI and that the occurrence of stroke predicts decreased activity and cognitive function.¹⁷ Given the association of BCVIs with traumatic brain injury, functional and cognitive impairment may be further exacerbated in the elderly in whom falls are common.

Although screening for BCVIs has traditionally been reserved for patients with high-energy mechanisms, recent studies have reported a higher-than-expected incidence in elderly patients (≥ 65) who were screened after low-energy mechanisms such as falls.^{18–19} In addition to BCVI being a known risk factor for ischemic strokes, elderly patients often have additional predisposing comorbidities such as atrial fibrillation, diabetes, hypertension, and atherosclerotic disease such as carotid artery stenosis.^{20–23} Given that the US population is aging rapidly,²⁰ there has been an increased focus on the management and outcomes of trauma patients who are older or have multiple comorbidities. However, little is known regarding the incidence of BCVI-related versus non-BCVI-related strokes and their outcomes, particularly in elderly patients.

The purpose of this study is to evaluate the incidence over time, etiology, risk factors, and in-hospital outcomes of strokes in a trauma population cared for at a high-volume level 1 trauma center. The primary hypothesis of the study is that elderly trauma patients are more likely to have non-BCVI-related strokes than BCVI strokes. The implication is that stroke prevention in trauma patients will require additional interventions beyond early diagnosis and treatment of BCVIs.

Figure 1.

Figure 1 Consort diagram.

Figure 2 Incidence of stroke over time.

METHODS

Study setting

The Red Duke Trauma Institute (RDTI) at Memorial Hermann Hospital (MHH) is an American College of Surgeons-verified Level 1 trauma center in Houston, Texas and serves as the primary teaching facility for McGovern Medical School at UT Houston. The RDTI treats over 7500 trauma patients per year and has approximately 5000 adult trauma admissions per year.

Selection of participants

After obtaining Institutional Review Board approval, a single-center retrospective study was performed using electronic medical records and the prospectively maintained trauma and stroke registries from MHH. All adult (≥ 16 years of age) trauma patients diagnosed with a stroke during the index trauma admission between January 2010 and December 2017 were included in this study.

Definitions

A BCVI-related stroke was defined as an ischemic stroke in the vascular distribution of the BCVI. Non-BCVI-related strokes were defined as ischemic strokes in the absence of BCVI or ischemic strokes in a vascular distribution not consistent with the BCVI. Adjudication was performed by a stroke neurologist not involved in the care of the patient who reviewed the imaging reports and studies. Time of CT angiography (CTA) neck acquisition was used as time of BCVI diagnosis. Time of treatment was the time of receipt of initial antithrombotic or

therapeutic anticoagulation medication. Time of stroke was the time of radiographic diagnosis of stroke with either CT or MRI brain. Death was classified as stroke-related based on divisional mortality review forms and independent clinical review.

The etiology of ischemic strokes were classified based on the Trial of ORG 10172 in Acute Stroke Treatment (TOAST) stroke subtype classification system.²⁴ The TOAST classification system classifies ischemic stroke based on the major pathophysiologic mechanisms and includes five categories: large artery atherosclerosis, cardioembolic, small vessel occlusion, stroke of other determined etiology, and stroke of undetermined etiology. Undetermined etiology includes patients with a negative evaluation, an incomplete evaluation, or with two or more potential causes identified. A vascular neurologist not involved in the care of the patient classified all of the strokes using the TOAST classification system.

Statistical analysis

A two-piece linear trend line was used to model the incidence rates of BCVI and strokes, including BCVI-related and non-BCVI-related strokes between 2011 and 2017. Significance for a trend was evaluated by the Wald test on zero slope, while the Jonckheere-Terpstra test was used to evaluate trends in age. Continuous characteristics were summarized as median and IQR. Association with type of stroke was evaluated by Wilcoxon rank-sum test and Fisher's exact test for continuous and categorical characteristics, respectively. Mortality rates and proportions of death due to stroke of BCVI and non-BCVI stroke were evaluated by both risk difference and relative risk. Relative risk is calculated as ratio of two proportions. We conducted the Wald tests to compare two proportions and test the relative risk equal to 1. We reported two-sided p values and p values less than 0.05 were considered as significant. All statistical analyses were performed using the SAS software (V.9.4; SAS Institute, Cary, NC).

RESULTS

Incidence

Of the 43 674 adult trauma patients admitted during the study period, 99 (0.2%) were diagnosed with an ischemic stroke during the index admission, of which only 21% were BCVI related (figure 1). The incidence of both BCVI and stroke, regardless of etiology, increased during the 7-year study period (figure 2). However, the incidence of BCVI-related stroke decreased while that of non-BCVI-related stroke increased (figure 3). The number of screening neck CTA studies also increased as a

Figure 3 Etiology of stroke over time.

Table 1 Baseline data for BCVI and non-BCVI stroke

	BCVI stroke Median, N (%) n=21	Non-BCVI stroke Median, N (%) n=78	P value
Age	44 (27 to 55)	71 (58 to 82)	<0.001
Male	7 (33.3%)	46 (59.0%)	0.078
Mechanism			<0.001
MVC	16 (76.2%)	22 (28.2%)	
MCC	0 (0%)	0 (0%)	
Auto-Ped	2 (9.5%)	2 (2.6%)	
Fall	2 (9.5%)	43 (55.1%)	
Other/unknown	1 (4.8%)	11 (14.1%)	
AIS head	3 (2 to 4)	0 (0 to 3)	<0.001
AIS face	0 (0 to 1)	0 (0 to 0)	0.003
AIS chest	2 (0 to 4)	0 (0 to 3)	0.009
AIS abdomen	0 (0 to 3)	0 (0 to 0)	0.005
AIS extremity	2 (0 to 3)	2 (0 to 3)	0.96
ISS	26 (20 to 34)	11 (8 to 18)	<0.001
BCVI grade*			<0.001
Grade 1	3 (14.3%)	1 (1.3%)	
Grade 2	6 (28.5%)	0 (0%)	
Grade 3	3 (14.3%)	0 (0%)	
Grade 4	9 (42.9%)	3 (3.9%)	
Grade 5	0 (0%)	1 (1.3%)	
BCVI location			<0.001
Vertebral artery	10 (47.6%)	0 (0%)	
Internal carotid artery	11 (52.4%)	5 (6.4%)	
GCS in ED	8 (3 to 15)	14.5 (9 to 15)	0.028
SBP in ED (mm Hg)	124 (100 to 146)	145 (120 to 169)	0.044
HR in ED (bpm)	94 (84 to 116)	90 (79 to 110)	0.38
Admission laboratory values			
Platelets (k/mm ³)	263 (209 to 299)	219 (174.5 to 280.5)	0.062
Hematocrit (%)	12.6 (11.7 to 13.7)	12.5 (10.9 to 13.5)	0.31
TEG ACT (s)	113 (97 to 121)	113 (97 to 121)	
TEG MA (mm)	67 (66 to 71)	64.5 (58 to 69)	
Medical comorbidities			
Any medical comorbidity†	12 (57.1%)	66 (84.6%)	0.013
Atrial fibrillation	0 (0%)	14 (18.0%)	0.037
Stroke or TIA‡	0 (0%)	17 (21.8%)	0.020
CAD/MI, CHF, PVD, hyperlipidemia, and HTN	11 (52.4%)	60 (76.9%)	0.053
Pre-hospital ASA or Plavix	1 (4.8%)	12 (15.4%)	0.29
Pre-hospital oral anticoagulant	0 (0%)	10 (12.8%)	0.11

Summary statistics for continuous variables are median and IQR. Associations between characteristics and type of stroke were evaluated by using Wilcoxon rank-sum test for continuous variables and Fisher's exact test for categorical variables.

*Five patients in the non-BCVI stroke group had a BCVI present on imaging but not responsible for the ischemic stroke.

†Medical comorbidities include stroke, transient ischemic attack (TIA), atrial fibrillation, coronary artery disease (CAD), myocardial infarction (MI), congestive heart failure (CHF), hypertension (HTN), peripheral vascular disease (PVD), hyperlipidemia, cancer, deep venous thrombosis (DVT), pulmonary embolism (PE), dementia, diabetes, end-stage renal disease, morbid obesity, smoker.

‡Pre-existing condition.

ACT, activated clotting time; ED, emergency department; HR, heart rate; MA, maximum amplitude; MCC, motorcycle crash; MVC, motor vehicle crash; SBP, systolic blood pressure; TEG, thromboelastography.

result of revisions to our BCVI screening criteria with 12.8% of patients receiving screening neck CTA compared with 5.5% at the beginning of the study period. The median age of adult trauma patients also increased from 41 to 46. All of these trends were statistically significant ($p < 0.001$).

Etiology

The majority (78/99, 78%) of strokes were non-BCVI related. Using the TOAST classification system, the etiologies of these strokes included, in descending order of frequency,

Table 2 Outcomes data for BCVI and non-BCVI stroke

	BCVI stroke Median, N (%) n=21	Non-BCVI stroke Median, N (%) n=78	P value
Hospital LOS (days)	14 (7 to 20)	12 (6 to 20)	0.85
ICU LOS (days)	7 (4 to 10)	3 (1 to 13)	0.29
Ventilator days	4 (0 to 7)	1 (0 to 7)	0.31
Mortality	8 (38.1%)	17 (21.8%)	0.16
% death due to stroke	75%	52.9%	0.26
NIHSS*	10 (2 to 23)	11 (2 to 22)	0.082
Operation with general anesthesia	17 (81%)	41 (52.6%)	
Time (h)			
ED arrival to BCVI diagnosis (h)	1.4 (1 to 3.6)	n/a	
Diagnosis to BCVI treatment (h)	15 (4.3 to 30.6)	n/a	
ED arrival to BCVI treatment (h)	18.2 (6.5 to 36)	n/a	
BCVI diagnosis to stroke (h)	24 (0 to 90)	n/a	
ED arrival to stroke (h)	44 (6.5 to 92.3)	47.3 (7.4 to 120)	0.53
Discharge disposition			0.032
Dead	8 (38.1%)	17 (21.8%)	
Home	2 (9.5%)	14 (17.9%)	
Rehabilitation	7 (33.3%)	8 (10.3%)	
SNF/LTAC†	3 (14.3%)	30 (38.5%)	
Hospice	1 (4.8%)	8 (10.3%)	
Other	0 (0%)	1 (1.3%)	

Summary statistics for continuous variables are median and IQR. Associations between characteristics and type of stroke were evaluated by using Wilcoxon rank-sum test for continuous variables and Fisher's exact test for categorical variables. Wald test was used for comparing mortality rates of two types of stroke.

*National Institutes of Health Stroke Scale.

†Skilled nursing facility (SNF) and long-term acute care facility (LTAC) values in parentheses without % are quartiles. ED, emergency department; ICU, intensive care unit; LOS, length of stay; n/a, not available.

cardioembolic (41%), other determined etiology (26%), undetermined etiology (13%), large artery atherosclerosis (11%), and small vessel occlusion (9%).

Risk factors

The BCVI-related stroke group were significantly younger than the non-BCVI-related stroke group. They also had a significantly higher median injury severity score (ISS), higher abbreviated injury scale (AIS) head, and lower Glasgow Coma Scale (GCS) score in the emergency department (table 1). Motor vehicle collisions were the most common mechanism of injury in the BCVI-related stroke group while ground level fall predominated in the non-BCVI stroke group. While patients with BCVI-related stroke were more likely to suffer multiple traumatic injuries from MVC, non-BCVI strokes had more isolated extremity injuries from fall mechanism.

The non-BCVI-related stroke group was older and more frequently had pre-existing medical comorbidities, especially atrial fibrillation and prior stroke or transient ischemic attack. Twelve of the 32 patients with a cardioembolic stroke had pre-existing atrial fibrillation: five patients had a stroke present on admission, three suffered a stroke while holding anticoagulants for surgery, two were taking no anticoagulants, and two suffered a stroke despite continuing aspirin. More patients with non-BCVI-related stroke were taking antiplatelet or anticoagulant medications, although this difference was not statistically significant.

Management of BCVIs

The median time from ED arrival to stroke diagnosis was similar for BCVI-related and non-BCVI-related strokes (table 2). Also, 95% of patients with BCVI received treatment with aspirin or heparin infusion. While the median time to diagnosis of BCVI

Box 1 Screening criteria* for BCVI

Injury patterns

- ▶ Complex facial fractures (LeFort II or III).
- ▶ Mandible fracture.
- ▶ Basilar skull fracture or occipital condyle fracture.
- ▶ Cervical vertebral body or transverse foramen fracture at any level (C1–7).
- ▶ Any fracture at level C1–C3.
- ▶ Cervical subluxation or ligamentous injury at any level.
- ▶ Severe traumatic brain injury (TBI) with GCS score <6.
- ▶ Near hanging with anoxic brain injury.
- ▶ Seatbelt or other clothesline-type injury with significant swelling, pain or AMS (Altered Mental Status).
- ▶ Combined TBI and major thoracic injury.
- ▶ Scalp degloving injury.
- ▶ Thoracic vascular injury.

Signs and symptoms

- ▶ Potential arterial hemorrhage from neck/nose/mouth.
- ▶ Cervical bruit in patient <50 years of age.
- ▶ Cervical hematoma.
- ▶ Focal neurologic defect: TIA, hemiparesis, vertebrobasilar symptoms, Horner's syndrome.
- ▶ Neurologic deficit inconsistent with head CT.
- ▶ Stroke on CT or MRI.

*The screening criteria are an adaptation of the expanded Denver criteria based on literature review and group consensus.

was only 1.4 hours, the median time of treatment initiation was 15 hours after the diagnosis. Eleven of the 21 patients with BCVI had a traumatic brain injury or solid-organ injury which contributed to the delay in starting antithrombotic therapy. Ischemic stroke was diagnosed in 10 patients prior to initiation of antithrombotic therapy (five patients with severe traumatic brain injury and five suffered a stroke prior to arrival to the ED) and 11 patients despite treatment of BCVI. Median time to stroke diagnosis was 44 hours after ED arrival with five patients diagnosed with stroke at time of arrival to the emergency center, resulting in a 24% non-preventable BCVI stroke rate.

Outcomes

Despite the non-BCVI-related stroke group having less severe injuries, the hospital length of stay, ICU length of stay, and number of ventilator days were similar to those in the BCVI-stroke group (table 2). The mortality rate for BCVI strokes and non-BCVI strokes were 38% and 22%, respectively, and patients with BCVI strokes were 1.7 times more likely to die in the hospital than those with non-BCVI strokes (relative risk (RR) 1.75, 95% CI 0.88 to 3.48, $p=0.11$). Of the patients who died, death was the direct result of the stroke more frequently in the BCVI group (75%) than the non-BCVI group (53%) (RR 1.42, 95% CI 0.78 to 2.58, $p=0.26$).

DISCUSSION

In this single-center study, ischemic strokes were rare with an incidence of 0.2% in an adult trauma inpatient population. Although low, the incidence has increased over the past 5 years, corresponding with an increasingly older population. The majority of strokes were not caused by BCVI but instead occurred in elderly patients with medical comorbidities and isolated extremity injuries from ground level fall. Only one of every five ischemic

strokes were BCVI related, and one out of four of those were present on admission and thus non-preventable. These findings have important implications on strategies to mitigate stroke risk.

For BCVI-related strokes, best practice includes following evidence-based screening guidelines for early BCVI diagnosis and treatment with antithrombotic therapy. At our institution, criteria for obtaining CTA to screen for BCVI have been in place since 2003 and were expanded in 2014 (box 1). As a result of the expansion of screening criteria, there was a significant increase in the number of CTA neck studies and a subsequent increase in BCVI diagnosis. The evidence-based management algorithm includes initiation of either aspirin or low-dose heparin infusion as soon as possible. Similar to the results reported by Shahan *et al.*,¹² this study demonstrates that aggressive screening for BCVI was associated with an increased incidence of BCVI. More importantly, there was a decreased risk of stroke among the entire trauma cohort. The increased BCVI incidence corresponds with changes to our evidence-based guidelines for diagnosis and treatment of BCVI.²⁵ Nonetheless, despite aggressive screening for BCVI, some strokes remained non-preventable such as those with stroke present on arrival or patients with severe traumatic brain injury and contraindication to immediate BCVI treatment.

For non-BCVI-related strokes, further studies are needed to identify high-risk elderly trauma patients and modifiable risk factors in order to focus efforts on stroke prevention. One potentially modifiable risk factor is occult hypoperfusion which is prevalent in the elderly population and associated with increased morbidity and mortality following trauma.^{26,27} Further study is necessary to determine the impact of early recognition and treatment of occult hypoperfusion on complications such as ischemic stroke.

The majority of strokes in this study were not BCVI related and occurred in patients with advanced age and medical comorbidities. The non-BCVI strokes occurred in elderly patients who were more likely to suffer an isolated extremity injury from a low mechanism of injury such as ground level fall. This observation has important implications regarding service admission patterns (non-surgical service vs trauma surgery) and the importance of optimization of these high-risk but low-acuity trauma patients. Regardless of admitting service, elderly patients may benefit from a comprehensive geriatric assessment, or an interdisciplinary approach to assessing and treating frail older patients' medical, psychosocial, and functional limitations.^{28,29} A recent systematic review of models of care for geriatric hip fracture patients demonstrated that patients under the care of a coordinated orthogeriatrics model or a geriatrics-led model had a shorter time to surgery, decreased length of stay, and reduced postoperative mortality rates.³⁰ A similar interdisciplinary approach has been demonstrated to reduce 30-day readmissions in a more diverse older trauma patient population.³¹ Whether such interdisciplinary care is achieved through geriatric consultation or development of a specialized geriatric trauma service with evidence-based protocols is dependent on each hospital's resources and surgeons' preferences.

This study has several limitations based on its retrospective design. Data were collected from the medical record, trauma registry, and stroke registry, which are only as accurate as documentation and data entry. The study did not include long-term follow-up data and included strokes occurring during the index hospital admission—not after discharge. Another limitation of the study is that there may be unmeasured confounders that affect the development of stroke.

Finally, since data were not collected on management of all patients with BCVI, the decreased stroke rate in patients

with BCVI cannot be directly attributed to the evidence-based management of these patients.

CONCLUSION

The incidence of stroke has increased over time at a large level 1 trauma center with aggressive BCVI screening and management protocols. The low number of BCVI-related strokes supports the practice of early identification and treatment of BCVIs. In addition, the increase in non-BCVI-related strokes in elderly patients with isolated extremity injuries secondary to ground level fall suggests the need to focus future efforts on managing medical comorbidities of an increasingly aging trauma population. Strategies to mitigate non-BCVI-related strokes should focus on optimizing medical conditions such as through early identification and treatment of occult hypoperfusion or through development of multidisciplinary geriatric protocols and services. Understanding of the incidence and etiology of strokes in trauma patients is necessary to guide future improvement efforts.

Contributors MKM, SIS, CEW, and LSK made substantial contributions to the conception and design of study. MKM, CS, DR, HKRI, XZ, BAC, and LSK made substantial contributions to the acquisition, analysis, and interpretation of the data. MKM wrote the manuscript. BAC, JH, CEW, SIS, and LSK performed critical revision of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The protocol was approved by the Institutional Review Board (IRB) at the University of Texas Health Science Center (HSC-GEN-13_0325). As this was a retrospective study, the consent by the subjects was not required by the IRB.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplementary information. Data will be made available under a Data Use Agreement through the University of Texas McGovern Medical School.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

John Harvin <http://orcid.org/0000-0002-2081-6256>

REFERENCES

- Biffi WL, Moore EE, Offner PJ, Brega KE, Franciose RJ, Elliott JP, Burch JM. Optimizing screening for blunt cerebrovascular injuries. *Am J Surg* 1999;178:517–21.
- Biffi WL, Moore EE, Ryu RK, Offner PJ, Novak Z, Coldwell DM, Franciose RJ, Burch JM. The unrecognized epidemic of blunt carotid arterial injuries: early diagnosis improves neurologic outcome. *Ann Surg* 1998;228:462–70.
- Burlew CC, Biffi WL, Moore EE, Barnett CC, Johnson JL, Bensard DD. Blunt cerebrovascular injuries. *J Trauma Acute Care Surg* 2012;72:330–7.
- Cothren CC, Moore EE, Ray CE, Johnson JL, Moore JB, Burch JM. Cervical spine fracture patterns mandating screening to rule out blunt cerebrovascular injury. *Surgery* 2007;141:76–82.
- Goodwin RB, Beery PR, Dorbish RJ, Betz JA, Hari JK, Opalek JM, MaGee DJ, Hinze SS, Scileppi RM, Franz RW, et al. Computed tomographic angiography versus conventional angiography for the diagnosis of blunt cerebrovascular injury in trauma patients. *J Trauma* 2009;67:1046–50.
- Eastman AL, Muraliraj V, Sperry JL, Minei JP. CTA-based screening reduces time to diagnosis and stroke rate in blunt cervical vascular injury. *J Trauma* 2009;67:551–6.
- Emmett KP, Fabian TC, DiCocco JM, Zarzaur BL, Croce MA. Improving the screening criteria for blunt cerebrovascular injury: the appropriate role for computed tomography angiography. *J Trauma* 2011;70:1058–65. discussion 1063–1055.
- DiCocco JM, Emmett KP, Fabian TC, Zarzaur BL, Williams JS, Croce MA. Blunt cerebrovascular injury screening with 32-channel multidetector computed tomography: more slices still don't cut it. *Ann Surg* 2011;253:444–50.
- Laser A, Kufera JA, Bruns BR, Sliker CW, Tesoriero RB, Scalea TM, Stein DM. Initial screening test for blunt cerebrovascular injury: validity assessment of whole-body computed tomography. *Surgery* 2015;158:627–35.
- Roberts DJ, Chaubey VP, Zygun DA, Lorenzetti D, Faris PD, Ball CG, Kirkpatrick AW, James MT. Diagnostic accuracy of computed tomographic angiography for blunt cerebrovascular injury detection in trauma patients: a systematic review and meta-analysis. *Ann Surg* 2013;257:621–32.
- Paulus EM, Fabian TC, Savage SA, Zarzaur BL, Botta V, Dutton W, Croce MA. Blunt cerebrovascular injury screening with 64-channel multidetector computed tomography: more slices finally cut it. *J Trauma Acute Care Surg* 2014;76:279–85. discussion 284–275.
- Shahan CP, Croce MA, Fabian TC, Magnotti LJ. Impact of continuous evaluation of technology and therapy: 30 years of research reduces stroke and mortality from blunt cerebrovascular injury. *J Am Coll Surg* 2017;224:595–9.
- Cothren CC, Biffi WL, Moore EE, Kashuk JL, Johnson JL. Treatment for blunt cerebrovascular injuries: equivalence of anticoagulation and antiplatelet agents. *Arch Surg* 2009;144:685–90.
- Stein DM, Boswell S, Sliker CW, Lui FY, Scalea TM. Blunt cerebrovascular injuries: does treatment always matter? *J Trauma* 2009;66:132–44.
- Burlew CC, Biffi WL, Moore EE, Pieracci FM, Beauchamp KM, Stovall R, Wagenaar AE, Jurkovich GJ. Endovascular stenting is rarely necessary for the management of blunt cerebrovascular injuries. *J Am Coll Surg* 2014;218:1012–7.
- Hundersmarck D, Slooff W-BM, Homans JF, van der Vliet QMJ, Moayeri N, Hietbrink F, de Borst GJ, Öner FC, Muijs SPJ, Leenen LPH, et al. Blunt cerebrovascular injury: incidence and long-term follow-up. *Eur J Trauma Emerg Surg* 2019. [Epub ahead of print: 13 Jun 2019].
- Shahan CP, Stavely TC, Croce MA, Fabian TC, Magnotti LJ. Long-term functional outcomes after blunt cerebrovascular injury: a 20-year experience. *Am Surg* 2018;84:551–6.
- Flashburg E, Ong AW, Muller A, Sherwood A, Wilhelm S, Zavilla J, Martin AP, Castor L, Barbera SC, Reinhart R, et al. Fall downs should not fall out: blunt cerebrovascular injury in geriatric patients after low-energy trauma is common. *J Trauma Acute Care Surg* 2019;86:1010–4.
- Anto VP, Brown JB, Peitzman AB, Zuckerbraun BS, Neal MD, Watson G, Forsythe R, Billiar TR, Sperry JL. Blunt cerebrovascular injury in elderly fall patients: are we screening enough? *World J Emerg Surg* 2018;13:30.
- Keller JM, Sciadini MF, Sinclair E, O'Toole RV. Geriatric trauma: demographics, injuries, and mortality. *J Orthop Trauma* 2012;26:e161–5.
- Adams SD, Holcomb JB. Geriatric trauma. *Curr Opin Crit Care* 2015;21:1–526.
- Hashmi A, Ibrahim-Zada I, Rhee P, Aziz H, Fain MJ, Friese RS, Joseph B. Predictors of mortality in geriatric trauma patients: a systematic review and meta-analysis. *J Trauma Acute Care Surg* 2014;76:894–901.
- Shaleah Levant KC, DeFrances C. Hospitalizations for patients aged 85 and over in the United States, 2000–2010. *National Center for Health Statistics Data Brief* 2015;182.
- Adams HP, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, Marsh EE. Classification of subtype of acute ischemic stroke. definitions for use in a multicenter clinical trial. TOAST. trial of ORG 10172 in acute stroke treatment. *Stroke* 1993;24:35–41.
- Division of Acute Care Surgery MMS. Screening for blunt cerebrovascular injury (BCVI) clinical practice policy. 2014.
- Neville AL, Nemtsev D, Manasrah R, Bricker SD, Putnam BA. Mortality risk stratification in elderly trauma patients based on initial arterial lactate and base deficit levels. *Am Surg* 2011;77:1337–41.
- Blow O, Magliore L, Claridge JA, Butler K, Young JS. The golden hour and the silver day: detection and correction of occult hypoperfusion within 24 hours improves outcome from major trauma. *J Trauma* 1999;47:964–9.
- Eamer G, Taheri A, Chen SS, Daviduck Q, Chambers T, Shi X, Khadaroo RG. Comprehensive geriatric assessment for older people admitted to a surgical service. *Cochrane Database Syst Rev* 2018;1:Cd012485.
- Trauma ACoSTQIPCo. ACS TQIP geriatric trauma management guidelines. 2013.
- Patel JN, Klein DS, Sreekumar S, Liporace FA, Yoon RS. Outcomes in multidisciplinary team-based approach in geriatric hip fracture care: a systematic review. *J Am Acad Orthop Surg* 2019. [Epub ahead of print: 30 May 2019].
- Bryant EA, Tulebaev S, Castillo-Angeles M, Moberg E, Senglaub SS, O'Mara L, McDonald M, Salim A, Cooper Z. Frailty identification and care pathway: an interdisciplinary approach to care for older trauma patients. *J Am Coll Surg* 2019;228:852–9.