


Participant retention in trauma intensive care unit (ICU) follow-up studies: a post-hoc analysis of a previous scoping review

Himanshu Rawal,¹ Daniel L Young,^{2,3,4} Roozbeh Nikooie,⁵ Awsse H Al Ani,⁶ Lisa Aronson Friedman,^{4,7} Sumana Vasishta,⁸ Elliott R Haut ,^{9,10,11,12,13} Elizabeth Colantuoni,^{4,14} Dale M Needham,^{3,4,7,15} Victor D Dinglas^{4,7}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/tsaco-2020-000584>).

For numbered affiliations see end of article.

Correspondence to

Victor D Dinglas; victor.dinglas@jhmi.edu

Received 15 August 2020
Revised 24 September 2020
Accepted 14 October 2020

ABSTRACT

Background The study aimed to synthesize participant retention-related data for longitudinal follow-up studies of survivors from trauma intensive care units (ICUs).

Methods Within a published scoping review evaluating ICU patient outcomes after hospital discharge, two screeners independently searched for trauma ICU survivorship studies.

Results There were 11 trauma ICU follow-up studies, all of which were cohort studies. Twelve months (range: 1–60 months) was the most frequent follow-up time point for assessment (63% of studies). Retention rates ranged from 54% to 94% across time points and could not be calculated for two studies (18%). Pooled retention rates at 3, 6, and 12 months were 75%, 81%, and 81%, respectively. Mean patient age (OR 0.85 per 1-year increase, 95% CI 0.73 to 0.99, $p=0.036$), percent of men (OR 1.07, 95% CI 1.04 to 1.10, $p=0.002$), and publication year (OR 0.89 per 1-year increase, 95% CI 0.82 to 0.95, $p=0.007$) were associated with retention rates. Early (3-month) versus later (6-month, 12-month) follow-up time point was not associated with retention rates.

Discussion Pooled retention rates were >75%, at 3-month, 6-month, and 12-month time points, with wide variability across studies and time points. There was little consistency with reporting participant retention methodology and related data. More detailed reporting guidelines, with better author adherence, will help improve reporting of participant retention data. Utilization of existing research resources may help improve participant retention.

Level of evidence Level III: meta-analyses (post-hoc analyses) of a prior scoping review.

INTRODUCTION

Advancement in trauma care has improved survival among critically injured patients, who often experience prolonged admissions in an intensive care unit (ICU).¹ Survivors often have reduced health-related quality of life.^{2–4} Hence, there is an increasing number of studies evaluating patient outcomes after hospital discharge, including survivors from trauma^{5,6} and trauma ICUs.⁷ This approach is critical to understanding the full reintegration of injured patients into society, as promoted by the National Academies of Sciences, Engineering, and Medicine report titled “A National Trauma Care System:

Integrating Military and Civilian Trauma Systems to Achieve Zero Preventable Deaths After Injury”^{8,9} and the National Quality Forum’s Population-Based Trauma Outcomes report.¹⁰

Retaining study participants in longitudinal follow-up studies can be challenging, but is integral for study validity and statistical power.¹¹ There is growing interest in understanding and implementing the most effective participant retention strategies^{12–14}; however, to our knowledge, there has been no synthesis of participant retention-related data across studies evaluating postdischarge outcomes of trauma ICU survivors. Thus, the objective of this article was to synthesize retention rates and strategies from studies evaluating postdischarge outcomes of trauma ICU survivors.

METHODS

This article follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses checklist.¹⁵ This systematic review’s protocol was registered with PROSPERO (International Prospective Register of Systematic Reviews; CRD42018087835).

Search strategy and study selection

The database from a prior comprehensive scoping review of 425 ICU survivorship articles that included at least one posthospital outcome measure was searched for publications on trauma ICU studies for inclusion in this analysis.⁷ The detailed search strategy and methods for this scoping review are reported elsewhere.⁷ In summary, the scoping review retrieved 20 189 citations from searching five online publication databases (PubMed, EMBASE, PsycINFO, Cumulative Index of Nursing and Allied Health Literature, and the Cochrane Controlled Trials Registry) during the designated search period (1970–2013). No language restrictions were applied in the scoping review.

From among the 425 articles reported in the scoping review, two trained researchers (HR and RN) independently screened full articles for studies that included trauma ICU patients. The researchers were not blinded to author/journal details. Studies were excluded if (1) non-trauma patients were included in the study or (2) there was only a single follow-up time point at which both consent and

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Rawal H, Young DL, Nikooie R, et al. *Trauma Surg Acute Care Open* 2020;**5**:e000584.

follow-up data collection occurred (ie, no prospective follow-up happened after consent).

Data abstraction

Duplicate data abstraction was performed by pairs of researchers. Conflicts were resolved by consensus, in consultation with a senior researcher (VDD or DMN). The following data were collected: participant retention rates and related data at each follow-up time point; reasons for loss to follow-up; use of a participant flow diagram; modes of data collection (eg, in person, phone, mail); reporting of mortality during follow-up; blinding of assessors (if interventional study); accounting for loss to follow-up in sample size/power calculation; study exclusion criteria related to barriers to follow-up (eg, homelessness); any discrepancy in reporting participant retention-related data; and description of participant retention strategies. Authors were contacted for additional data when necessary.

Risk of bias

There were no randomized controlled trials included in this systematic review. For observational studies, risk of bias was assessed using a modified Newcastle-Ottawa Scale,¹⁶ excluding three criteria not applicable to this systematic review given its focus on participant retention rather than a specific clinical end point: (1) demonstration that the outcome was not present at enrollment, (2) assessment of the outcome and (3) follow-up long enough for the outcome to occur.

Statistical analysis

Pooled average participant retention rates were calculated in this analysis. Among eligible studies, the following follow-up time points were reported: 3, 6, 12, 24, 36, and 60 months. Data from follow-up times >12 or <3 months could not be pooled due to only a single study evaluating the time point. For studies reporting participant age as median and IQR, for purposes of this analysis, mean and SD for participant age were estimated using the methods proposed by Wan *et al.*¹⁷ One study¹⁸ published mean and SD age separately for four treatment groups; these data, along with the sample size in each group, were used to calculate an overall mean and SD age for the study. One study supplied CI for age instead of SD, and the CI was converted to SD. Participant retention data reported separately for treatment groups or patient subgroups at the same time point within a study were tested for a statistically significant difference using Fisher's exact test and grouped if not significant. For studies where retention rates were 100%, the Haldane-Anscombe correction was used to correct the CI.^{19 20}

Participant retention rates were calculated in two ways. For the primary approach, the retention rate was calculated by dividing the number of participants who had a study assessment (numerator) by the total number of participants alive and eligible for follow-up at that same time point (denominator). The secondary definition also excluded those who withdrew from the study in calculating the denominator. Retention rates were not calculated if all requisite data were not reported or if mortality was combined with loss to follow-up data.

A linear random intercept regression model (logit transformation) was used to pool retention rates across all eligible studies and time points, where each study was represented by a value of the random intercept. This regression model was extended to determine if the pooled average retention rate was associated with two patient demographic characteristics reported in all studies (average age and percent of men) and with study




Figure 1 Flow diagram of identification of eligible studies on trauma ICU survivors. ICU, intensive care unit.

publication year. A separate extended regression model was constructed for each of the patient and study characteristics.

Statistical heterogeneity among included studies was evaluated using the I^2 statistics (with >50% deemed to be substantial heterogeneity).²¹ The I^2 statistics were calculated for each time point when there were more than >2 studies reporting data.²² SAS V.9.4 was used to conduct all analyses.

RESULTS

Of the 425 publications included in the original scoping review, 16 publications, reporting on 11 unique studies (figure 1), met the eligibility criteria for this analysis (ie, focused on trauma ICU survivors). All 11 studies were cohort studies.^{18 23–36 37} The time points for follow-up ranged from 1 to 60 months, with the most frequent time point being 12 months, occurring in seven (63%)^{25–27 29 31 33 36 37} studies (table 1). Five (45%) studies were exclusively conducted in Europe and four (36%) in the USA (table 1). In each of the 11 eligible studies, a majority of the participants were male, with a range of 57% to 83%. Across studies, the mean age ranged from 27 to 44 years old.

Risk of bias assessment

Of the 11 cohort studies, 6 had adequate reporting of follow-up (online supplemental table S1). Nine studies had a positive rating for comparability of cohorts. Lastly, two studies compared their trauma ICU cohort with a non-trauma cohort.

Reporting of retention data

Eight (72%) studies^{23–25 27 29 31 33 35–37} reported exclusion criteria related to ability to follow up participants after hospital discharge, with the most common exclusion criterion being language proficiency in five (62%) studies^{23 25 27 29 31} (table 2). No study reported accounting for loss to follow-up to calculate sample size or statistical power. Nine (82%) studies^{18 23–27 29 31 33 34 37} reported loss to follow-up and mortality data separately, with two (18%) publications^{35 36} combining them in their study reporting. A flow diagram for patient follow-up was included

Table 1 Study cohort characteristics

Publications for each unique study	Start and end of study	Type of study	Countries	Sample size*	Age† (years), mean (SD)	Male (%)	Modes of contact ever used in follow-up?	Time points for follow-up (months)	Retention rate Primary definition‡(%)	Retention rate Secondary definition§ (%)
Aitken <i>et al</i> ²³	June 2008–August 2012	Cohort	Australia	112	40 (20)	83	Phone, mail	1	76	83
Christensen <i>et al</i> ¹⁸	August 2005–December 2008	Cohort	USA and 26 countries	105	38 (14)	77	Not reported	6	72	84
Richards <i>et al</i> ^{24,25,27}	July 2006–July 2008	Cohort	USA	472	43 (17)	57	In person	3	72	74
Orwellius <i>et al</i> ²⁶	August 2000–June 2006	Cohort	Sweden	160	44 (18)	68	Mail	12	67	68
Schnyder <i>et al</i> ^{27,28}	January 1996–June 1998	Cohort	Switzerland	108	38 (13)	74	Not reported	12	79	79
Tøien <i>et al</i> ^{29,30}	June 2005–December 2007	Cohort	Norway	83	40 (16)	70	Mail	24	54	69
Hepp <i>et al</i> ^{31,32}	January 1996–June 2003	Cohort	Switzerland	121	38 (13)	74	Not reported	12	88	88
Davdow <i>et al</i> ³³	July 2001–September 2003	Cohort	USA	106	41 (32)	71	Phone	3	79	80
Frutiger <i>et al</i> ³⁴	1980–1988	Cohort	Switzerland	147	36 (17)	81	In person, other	12	79	80
Holbrook <i>et al</i> ³⁵	January 1990–June 1990	Cohort	USA	121	30 (13)	74	Phone	6	88	88
Mackenzie <i>et al</i> ³⁶	July 1982–March 1984	Cohort	USA	90	27 (7)	78	In person, phone	36	74	100
				1781	41 (32)	71	Phone	12	83	83
				177	36 (17)	81	In person, other	60	94	94
				–	30 (13)	74	Phone	3	NA¶	NA¶
				–	27 (7)	78	In person, phone	6	NA	NA
				–	–	–	–	12	NA	NA

*Sample size was calculated as the number of ICU survivors eligible for follow-up at hospital discharge. For studies that did not provide this information we used the sample size at start of the study, and for studies that did not provide either we used the sample size after informed consent was obtained. In cases where there were articles published while the study was still ongoing, we used the article with the largest sample size.

†Age was expressed as mean (SD) or median (IQR).

‡Primary participant retention rates were calculated as the number of participants assessed at each follow-up time point divided by the number presumed alive and eligible for assessment at that time point (this included the participants who withdrew and withdrawn just prior to the time point).

§Secondary participant retention rates were calculated after a sensitivity analysis where we calculated the number of participants assessed at each follow-up time point divided by the number alive and eligible for assessment at the time point (this *did not* include the participants who died, withdrew, and withdrawn just prior to the time point).

¶Retention rates could not be calculated because mortality was combined with loss to follow-up.

ICU, intensive care unit; NA, not applicable.

Table 2 Participant retention-related data in longitudinal studies of trauma ICU survivors

Participant retention-related issue	Studies reporting, n (%) (N=11)
Study exclusion criteria included barrier(s) to follow-up (eg, homelessness)	8 (72)
Sample size or power calculation	0 (0)
Reported use of strategies to improve participant retention	4 (36)
Mortality reported during follow-up	8 (72)
Reported lost to follow-up rates combined with mortality	2 (18)
Included flow diagram with retention rate for each follow-up time point	5 (45)
Reported reasons for lost to follow-up at each follow-up time point	4 (36)


ICU, intensive care unit.

in five (45%) studies, with only four (36%)^{26 27 31 34} reporting reasons for lost to follow-up at each time point.

Participant retention

In the nine studies with adequate reporting, retention rates could be calculated for all time points (range: 54%–94%). Pooled average retention rates (95% CI) at 3-month, 6-month, and 12-month follow-up time points were 75% (22% to 97%; 2 studies; n=632), 81% (1% to 100%; 2 studies; n=244), and 81% (72% to 88%; 6 studies; n=2441; $I^2=86\%$), respectively (figure 2).

Retention rates of the earliest (3 months) time point were not statistically different from the later time points at 6 months ($p=0.653$) or 12 months ($p=0.278$). For every 1-year increase in average participant age in the eligible studies, the odds of retention were lower by 15% (OR 0.85, 95% CI 0.73 to 0.99, $p=0.036$). For every 1% increase in the proportion of male participants in the eligible studies, the odds of retention were higher by 7% (OR 1.07, 95% CI 1.04 to 1.10, $p=0.002$). Finally,



N studies	2	2	6
N participants	632	244	2441
Average cohort retention rate	75	81	81

Figure 2 Pooled average retention rates in trauma ICU survivor follow-up studies. Retention rates were calculated as the number of participants assessed at each follow-up time point divided by the number presumed alive at that time point (this included the participants who withdrew and withdrawn just prior to the time point). Diamonds in the graph are the pooled average retention rates, whereas bars represent 95% CI. Linear random effects regression model was used to pool retention rates across all eligible studies and time points. ICU, intensive care unit.

publication year was also significantly associated with retention rate; with every 1-year increase (ie, 1-year more recent publication), the odds of retention were lower by 11% (OR 0.89, 95% CI 0.82 to 0.95, $p=0.007$). These results did not qualitatively change when evaluating the participant retention rates using the secondary definition, as previously described in the Methods section.

DISCUSSION

In this analysis, we report and synthesize participant retention-related data for 11 longitudinal studies reporting on functional outcomes of adult trauma ICU survivors. Two (18%) of the studies did not report adequate data for calculating retention rates. Among the remaining nine studies, retention rates ranged from 54% to 94% across follow-up time points, with pooled retention rates at 3, 6, and 12 months of 75%, 81%, and 81%, respectively.

The pooled retention rates from this analysis (75%–81%, during the first year of follow-up) were similar to posthospital follow-up rates from 21 studies of acute respiratory failure survivors (82%–89% during the first 2 years of follow-up).³⁸ These findings were also similar to a broader range of predominantly non-trauma/non-critical illness healthcare-related follow-up studies, as reported in a systematic review of 82 studies that reported retention strategies and rates (median 85%, IQR 79%–92%).^{12 13}

Timing of follow-up (3 months vs. 6 months or 12 months) was not associated with a difference in retention rates, but mean age, proportion of male participants, and publication year of studies were significantly associated with retention rates. Retention rates were higher with a greater proportion of male participants, and lower with older participants and with more recent study publication.

The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guidelines were published in 2007.³⁹ STROBE recommends reporting the number of “potentially eligible” participants and “confirmed eligible” participants, reporting reasons for non-participation, and including a participant flow diagram. Participant retention rates were reported in 100% of studies published after 2007, but only 60% studies before 2007. In addition, the two studies^{35 36} for which retention rates could not be calculated were published before the STROBE guidelines. Hence, perhaps STROBE guidelines have helped improve reporting in studies of trauma ICU survivors.

No study reported a sample size or statistical power calculation, and <50% of studies reported other important research methodology components (eg, participant retention strategies used, participant flow diagram). Reporting on loss to follow-up data varied widely. We were unable to calculate retention rates in two (18%) studies because loss to follow-up was combined with mortality. These findings highlight the potential value of updating the STROBE guidelines to require more detailed reporting. Interestingly, the majority of the studies that reported retention strategies had high retention rates.

To reduce selection bias in follow-up studies, there is growing interest in participant retention and related methodology, as evidenced by an increase in publications and resources focused on improving participant retention.¹³ The National Institutes of Health/National Heart, Lung, and Blood Institute funded a national research infrastructure project (R24HL111895), with one aim specifically focused on improving participant retention via creation and dissemination of practical retention

tools and resources to aid investigators (www.improvelto.com/cohort-retention-tools/).

For example, this project supported completion of a systematic review on participant retention strategies.^{12–13} From the studies included in that systematic review, the project compiled 618 participant retention strategies, across 12 different themes, which are available as a free searchable online database (www.improvelto.com/sysrevstrategies). Moreover, best practices for participant retention in healthcare-related studies have been published,¹⁴ along with four empirical analyses relating to participant retention.^{40–43} Such publications are important in ensuring evidence-based advancement of methods for participant retention.

Furthermore, one of the publications from this project provides empirical evidence to debunk the myth that intensive retention efforts are bothersome to participants.⁴¹ Ultimately, this national infrastructure project has shared >30 downloadable tools, including customizable telephone scripts and letters, as well as templates relevant to participant follow-up, such as a detailed participant contact information form. With increasing interest in posthospital outcomes of trauma patients,^{5–10–44} improving participant retention in studies evaluating long-term outcomes is critical to help reduce bias and better inform the care of critically injured patients.

Strengths and limitations

To our knowledge, this is the first evaluation of participant retention methodology in studies of adult trauma ICU survivors. There are potential limitations to be acknowledged. First, there are a relatively small number of studies, and studies published after 2013 could not be included since that was the end date of the database of studies from the prior scoping review on which this analysis was based. Second, there is heterogeneity in the studies that were pooled; hence, caution is advised in interpreting the pooled average retention rates, with recognition that there is some variability across studies and time points. Third, other factors that may be relevant to retention of post-ICU patients, such as discharge location, were not collected in this synthesis and should be considered in future studies. Lastly, since the focus of this analysis was adult trauma ICU survivors, these results may not generalize to other populations of critically ill patients.

CONCLUSION

In this evaluation of 11 studies of trauma ICU survivors, the pooled participant retention rate was >75% across 3-month, 6-month, and 12-month follow-up assessments. However, retention rates across individual studies were highly variable (54%–94%) and there was inconsistent reporting of retention-related methodological data. Although guidelines (Consolidated Standards of Reporting Trials and STROBE) recommend reporting participant retention data, more detailed guidance on such reporting, along with strict adherence by researchers, may help further advance research aiming to understand the postdischarge outcomes of trauma ICU survivors. Moreover, use of existing participant retention resources, including new NIH-funded free resources (see www.improveLTO.com), may help researchers mitigate loss to follow-up and its associated potential for low statistical power and bias.

Author affiliations

¹Pulmonary Disease and Critical Care Medicine, Wake Forest School of Medicine, Winston-Salem, North Carolina, USA

²Department of Physical Therapy, University of Nevada Las Vegas, Las Vegas, Nevada, USA

³Department of Physical Medicine and Rehabilitation, Johns Hopkins Medicine School of Medicine, Baltimore, Maryland, USA

⁴Outcomes After Critical Illness and Surgery Group, Johns Hopkins University, Baltimore, Maryland, USA

⁵Department of Internal Medicine, Yale-New Haven Hospital, New Haven, Connecticut, USA

⁶MedStar Union Memorial Hospital, Baltimore, Maryland, USA

⁷PCCM, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

⁸Institute of Nephro Urology Mysuru Branch, Krishna Rajendra Hospital Campus, Mysuru, India

⁹Division of Acute Care Surgery, Department of Surgery, School of Medicine, Johns Hopkins University, Baltimore, MD, United States

¹⁰Department of Anesthesiology and Critical Care Medicine, School of Medicine, Johns Hopkins University, Baltimore, MD, United States

¹¹Department of Emergency Medicine, School of Medicine, Johns Hopkins University, Baltimore, MD, United States

¹²The Armstrong Institute for Patient Safety and Quality, Johns Hopkins Medicine, Baltimore, MD, United States

¹³Department of Health Policy and Management, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, United States

¹⁴Department of Biostatistics, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland, USA

¹⁵School of Nursing, Johns Hopkins University, Baltimore, MD, United States

Contributors VDD and DMN contributed to conception and design of the article. HR, DLY, RN, AHAA, LAF, SV, ERH, EC, DMN, and VDD contributed to analysis and interpretation of data. HR and DLY drafted the article and all other authors critically revised it for important intellectual content. All authors gave final approval of the article version to be published. All authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding This research was supported by the National Heart, Lung, and Blood Institute (R24HL111895).

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Elliott R Haut <http://orcid.org/0000-0001-7075-771X>

REFERENCES

- MacKenzie EJ, Rivara FP, Jurkovich GJ, Nathens AB, Frey KP, Egleston BL, Salkever DS, Scharfstein DO. A national evaluation of the effect of trauma-center care on mortality. *N Engl J Med* 2006;354:366–78.
- Sluys K, Häggmark T, Iselius L. Outcome and quality of life 5 years after major trauma. *J Trauma* 2005;59:223–32.
- Ulvik A, Kvåle R, Wentzel-Larsen T, Flaatten H. Quality of life 2–7 years after major trauma. *Acta Anaesthesiol Scand* 2008;52:195–201.
- Eddleston JM, White P, Guthrie E. Survival, morbidity, and quality of life after discharge from intensive care. *Crit Care Med* 2000;28:2293–9.
- Haider AH, Herrera-Escobar JP, Al Rafai SS, Harlow AF, Apoj M, Nehra D, Kasotakis G, Brasel K, Kaafarani HMA, Velmahos G, et al. Factors associated with long-term outcomes after injury: results of the functional outcomes and recovery after trauma emergencies (Forte) multicenter cohort study. *Ann Surg* 2020;271:1165–73.
- Rios-Diaz AJ, Herrera-Escobar JP, Lilley EJ, Appelson JR, Gabbe B, Brasel K, deRoos-Cassini T, Schneider EB, Kasotakis G, Kaafarani H, et al. Routine inclusion of long-term functional and patient-reported outcomes into trauma registries: the Forte project. *J Trauma Acute Care Surg* 2017;83:97–104.
- Turnbull AE, Rabiee A, Davis WE, Nasser MF, Venna VR, Lolitha R, Hopkins RO, Bienvenu OJ, Robinson KA, Needham DM, et al. Outcome measurement in ICU survivorship research from 1970 to 2013: a scoping review of 425 publications. *Crit Care Med* 2016;44:1267–77.
- National Academies of Sciences, Engineering, and Medicine. *A national trauma care system: integrating military and civilian trauma systems to achieve zero preventable deaths after injury*. The National Academies Press, 2016.
- Haut E, Mann N, Kotwal R. *Military Trauma Care's Learning Health System: The Importance of Data Driven Decision Making*. Washington DC: The National Academies Press, 2016.

- 10 National Quality Forum. *Population-Based trauma outcomes*. Washington DC, 2019:1–69.
- 11 Fewtrell MS, Kennedy K, Singhal A, Martin RM, Ness A, Hadders-Algra M, Koletzko B, Lucas A. How much loss to follow-up is acceptable in long-term randomised trials and prospective studies? *Arch Dis Child* 2008;93:458–61.
- 12 Robinson KA, Dennison CR, Wayman DM, Pronovost PJ, Needham DM. Systematic review identifies number of strategies important for retaining study participants. *J Clin Epidemiol* 2007;60:757.e1–757.e19.
- 13 Robinson KA, Dinglas VD, Sukrithan V, Yalamanchilli R, Mendez-Tellez PA, Dennison-Himmelfarb C, Needham DM. Updated systematic review identifies substantial number of retention strategies: using more strategies retains more study participants. *J Clin Epidemiol* 2015;68:1481–7.
- 14 Abshire M, Dinglas VD, Cajita MA, Eakin MN, Needham DM, Himmelfarb CD. Participant retention practices in longitudinal clinical research studies with high retention rates. *BMC Med Res Methodol* 2017;17:1471–2288. (Linking):30.
- 15 Moher D, Liberati A, Tetzlaff J, Altman DG, . PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–9.
- 16 Wells G, Shea B, O'connell D. *The Newcastle-Ottawa scale (NOS) for assessing the quality of Nonrandomised studies in meta-analyses*: Ottawa Hospital Research Institute, 2000. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.
- 17 Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014;14:135.
- 18 Christensen MC, Banner C, Lefering R, Vallejo-Torres L, Morris S. Quality of life after severe trauma: results from the global trauma trial with recombinant factor VII. *J Trauma* 2011;70:1524–31.
- 19 Haldane JBS. The mean and variance of χ^2 , when used as a test of homogeneity, when expectations are small. *Biometrika* 1940;31:346–55.
- 20 Anscombe FJ. On estimating binomial response relations. *Biometrika* 1956;43:461–4.
- 21 Sedgwick P. Meta-Analyses: heterogeneity and subgroup analysis. *BMJ* 2013;346:f4040.
- 22 Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- 23 Aitken LM, Chaboyer W, Schuetz M, Joyce C, Macfarlane B. Health status of critically ill trauma patients. *J Clin Nurs* 2014;23:704–15.
- 24 Richards JE, Guillaumondegui OD, Archer KR, Jackson JC, Ely EW, Obremsky WT. The association of reamed intramedullary nailing and long-term cognitive impairment. *J Orthop Trauma* 2011;25:707–13.
- 25 Guillaumondegui OD, Richards JE, Ely EW, Jackson JC, Archer KR, Archer-Swygert K, Norris PR, Obremsky WT. Does hypoxia affect intensive care unit delirium or long-term cognitive impairment after multiple trauma without intracranial hemorrhage? *J Trauma* 2011;70:910–5.
- 26 Orwelius L, Bergkvist M, Nordlund A, Simonsson E, Nordlund P, Bäckman C, Sjöberg F. Physical effects of trauma and the psychological consequences of preexisting diseases account for a significant portion of the health-related quality of life patterns of former trauma patients. *J Trauma Acute Care Surg* 2012;72:504–12.
- 27 Schnyder U, Moergeli H, Klaghofer R, Buddeberg C. Incidence and prediction of posttraumatic stress disorder symptoms in severely injured accident victims. *Am J Psychiatry* 2001;158:594–9.
- 28 Schnyder U, Moergeli H, Trentz O, Klaghofer R, Buddeberg C. Prediction of psychiatric morbidity in severely injured accident victims at one-year follow-up. *Am J Respir Crit Care Med* 2001;164:653–6.
- 29 Tøien K, Bredal IS, Skogstad L, Myhren H, Ekeberg O. Health related quality of life in trauma patients. data from a one-year follow up study compared with the general population. *Scand J Trauma Resusc Emerg Med* 2011;19:22.
- 30 Tøien K, Myhren H, Bredal IS, Skogstad L, Sandvik L, Ekeberg Øivind, Ekeberg O. Psychological distress after severe trauma: a prospective 1-year follow-up study of a trauma intensive care unit population. *J Trauma* 2010;69:1552–9.
- 31 Hepp U, Moergeli H, Büchi S, Bruchhaus-Steinert H, Kraemer B, Sensky T, Schnyder U. Post-Traumatic stress disorder in serious accidental injury: 3-year follow-up study. *Br J Psychiatry* 2008;192:376–83.
- 32 Hepp U, Moergeli H, Büchi S, Wittmann L, Schnyder U. Coping with serious accidental injury: a one-year follow-up study. *Psychother Psychosom* 2005;74:379–86.
- 33 Davydov DS, Zatzick DF, Rivara FP, Jurkovich GJ, Wang J, Roy-Byrne PP, Katon WJ, Hough CL, Kross EK, Fan M-Y, et al. Predictors of posttraumatic stress disorder and return to usual major activity in traumatically injured intensive care unit survivors. *Gen Hosp Psychiatry* 2009;31:428–35.
- 34 Frutiger A, Ryf C, Bilal C, Rosso R, Furrer M, Cantieni R, Rüedi T, Leutenegger A. Five years' follow-up of severely injured ICU patients. *J Trauma* 1991;31:1216–26. discussion 1225–1226.
- 35 Holbrook TL, Hoyt DB, Anderson JP, Hollingsworth-Fridlund P, Shackford SR. Functional limitation after major trauma: a more sensitive assessment using the Quality of Well-being scale—the trauma recovery pilot project. *J Trauma* 1994;36:74–8.
- 36 MacKenzie EJ, Siegel JH, Shapiro S, Moody M, Smith RT. Functional recovery and medical costs of trauma: an analysis by type and severity of injury. *J Trauma* 1988;28:281–97.
- 37 Jackson JC, Archer KR, Bauer R, Abraham CM, Song Y, Greevey R, Guillaumondegui O, Ely EW, Obremsky W. A prospective investigation of long-term cognitive impairment and psychological distress in moderately versus severely injured trauma intensive care unit survivors without intracranial hemorrhage. *J Trauma* 2011;71:860–6.
- 38 Nunna K, Al-Ani A, Nikooie R, Friedman LA, Raman V, Wadood Z, Vasishtha S, Colantuoni E, Needham DM, Dinglas VD, et al. Participant retention in follow-up studies of acute respiratory failure survivors. *Respir Care* 2020;65:1382–91.
- 39 von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, . STROBE Initiative. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;370:1453–7.
- 40 Dinglas VD, Huang M, Sepulveda KA, Pinedo M, Hopkins RO, Colantuoni E, Needham DM, . NIH NHLBI ARDS Network. Personalized contact strategies and predictors of time to survey completion: analysis of two sequential randomized trials. *BMC Med Res Methodol* 2015;15:5.
- 41 Eakin MN, Eckmann T, Dinglas VD, Akinremi AA, Hosey M, Hopkins RO, Needham DM. Association between participant contact attempts and reports of being bothered in a national, longitudinal cohort study of ARDS survivors. *Chest* 2020;158:588–95.
- 42 Heins SE, Wozniak AW, Colantuoni E, Sepulveda KA, Mendez-Tellez PA, Dennison-Himmelfarb C, Needham DM, Dinglas VD. Factors associated with missed assessments in a 2-year longitudinal study of acute respiratory distress syndrome survivors. *BMC Med Res Methodol* 2018;18:55.
- 43 Heins SE, Wozniak AW, Colantuoni E. Factors associated with home visits in a 5-year study of acute respiratory distress syndrome survivors. *BMC Med Res Methodol* 2020;18:55.
- 44 Sakran JV, Ezzeddine H, Schwab CW, Bonne S, Brasel KJ, Burd RS, Cuschieri J, Ficke J, Gaines BA, Giacino JT, et al. Proceedings from the consensus conference on trauma patient-reported outcome measures. *J Am Coll Surg* 2020;230:819–35.