

Urine drug screen positive for cocaine and amphetamine is not an adverse risk factor for cardiovascular morbidity or mortality in trauma

Sangeeta Satish, Carl Freeman, John Culhane

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/tsaco-2021-000749>).

Trauma, Saint Louis University, Saint Louis, Missouri, USA

Correspondence to
Dr John Culhane; john.culhane@health.slu.edu

Received 24 March 2021
Accepted 3 August 2021

ABSTRACT

Background Urine drug screening (UDS) is a component of trauma workup and of perioperative risk evaluation. Illicit stimulant use has been associated with cardiovascular complications. This study investigates the impact of stimulant use and its interaction with surgery on cardiovascular complications in trauma patients.

Methods Patients were identified from the 2017 National Trauma Data Bank. Univariate and multivariate analyses were used to evaluate the effect of amphetamine and cocaine on mortality, myocardial infarction (MI), and stroke. We evaluated three subsets: all screened patients, those who underwent surgery, and those whose surgery was immediate. Significance was tested with χ^2 test for categorical variables, Student's t-test for continuous variables, and logistic regression for multivariate analysis.

Results 317 688 (32.1%) patients underwent UDS. Multivariate analysis showed protective association between cocaine and mortality OR 0.9 ($p=0.028$). Cocaine was a non-significant predictor of MI and stroke: OR 0.63 ($p=0.065$) and 0.91 ($p=0.502$), respectively. Amphetamine was a non-significant predictor of mortality, MI, and stroke: OR 0.97 ($p=0.405$), 0.80 ($p=0.283$), and 1.02 ($p=0.857$), respectively. On univariate analysis, amphetamine showed a protective association with MI for all screened patients: relative risk (RR) 0.58 ($p=0.005$), and for surgical patients: RR 0.58 ($p=0.019$). Amphetamine showed a protective association with mortality for all three subsets: RR 0.83 ($p<0.001$), 0.78 ($p<0.001$), and 0.71 ($p<0.001$), respectively. Cocaine showed a protective association with MI for all screened patients: RR 0.45 ($p=0.001$), and for surgical patients: RR 0.44 ($p=0.005$). Cocaine showed a protective association with mortality for all three subsets: RR 0.76 ($p<0.001$), 0.71 ($p<0.001$), and 0.63 ($p<0.001$), respectively.

Discussion UDS positive for cocaine or amphetamine is not an adverse risk factor in trauma, including trauma patients who underwent surgery. The apparent protective effects of illicit drugs warrant further investigation.

Level of evidence Therapeutic/care management, level IV.

INTRODUCTION

Discretionary urine drug screening (UDS) is a common component of a trauma workup. It has been shown to have a prognostic value which can help clinicians allocate resources and monitor patients with the appropriate acuity.¹ UDS is also frequently performed as part of a preoperative

evaluation for elective surgery. A positive drug test for stimulants, such as cocaine or methamphetamine, is generally perceived as an adverse perioperative cardiac risk factor resulting in cancellation of the procedure.² Stimulant use has been associated with an increased risk of cardiac complications, such as tachycardia, hypertension, arrhythmias and myocardial infarction (MI), due to overstimulation of the sympathetic nervous system, vasoconstriction and dysregulation of sodium/potassium channels in the heart.^{3,4} However, there is a lack of information on whether the risk of cardiac complications remains elevated in asymptomatic patients who do not show signs of stimulant toxicity. UDS can remain positive for days after drug use, which presents a challenge in evaluating perioperative risk.⁵

A patient population that routinely undergoes surgery despite having a positive drug screen is that of trauma patients, who often require emergent procedures that cannot be delayed. Trauma patients have a high prevalence of illicit substance use, with an estimated frequency of preinjury substance use of 20% to 50%.^{6,7} A recent study done by our group⁸ investigated the cardiac risk associated with surgical procedures in patients who had a positive UDS using the National Trauma Data Bank (NTDB). The results showed that a positive drug screen was not associated with increased perioperative cardiac morbidity or mortality, and in fact was a marker of lower risk. The analysis was done using an NTDB data set that only characterizes drug screen results as positive or negative, with no further breakdown into what substance was responsible for the positive test. Therefore, it is possible that the effect of cardioactive drugs was masked by other drugs responsible for the positive screen. Recently, the NTDB has provided more detailed information on the specific class of drug that led to a positive drug screen. Therefore, the goal of this study is to investigate the impact of stimulant use on perioperative mortality and cardiovascular risk in trauma patients. Our hypothesis is that a UDS positive for stimulant drugs is associated with increased cardiovascular risk and mortality in trauma patients, both for the operative and non-operative subsets of the population. The null hypothesis is that there is no association.

METHODS

The NTDB is a registry of trauma data from multiple US trauma centers. The year 2017 was chosen because this is the first year for which specific drugs

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Satish S, Freeman C, Culhane J. *Trauma Surg Acute Care Open* 2021;**6**:e000749.

Table 1 Baseline characteristics

Characteristics	No amphetamine	Amphetamine	No cocaine	Cocaine
Sex, female	96 986 (33.7%)	8505 (28.8%)	99 385 (33.9%)	6106 (24.6%)
Race (white vs. non-white)	195 290 (67.8%)	22 218 (75.2%)	203 241 (69.4%)	14 267 (57.5%)
Hispanic ethnicity	14 498 (28.5%)	3577 (45.6%)	14 564 (28.1%)	3511 (51.1%)
Angina pectoris	231 (0.1%)	9 (0.0%)	232 (0.1%)	8 (0.0%)
Myocardial infarction	2083 (0.7%)	157 (0.5%)	2115 (0.7%)	125 (0.5%)
Congestive heart failure	7855 (2.7%)	543 (1.8%)	7993 (2.7%)	405 (1.6%)
Hypertension	76 997 (26.7%)	5362 (18.2%)	77 528 (26.5%)	4831 (19.5%)
Diabetes mellitus	30 797 (10.7%)	2038 (6.9%)	30 990 (10.6%)	1845 (7.4%)
Peripheral arterial disease	1035 (0.4%)	67 (0.2%)	1043 (0.4%)	59 (0.2%)
Stroke	5786 (2.0%)	359 (1.9%)	5805 (2.0%)	340 (1.4%)
Chronic renal failure	2787 (1.0%)	120 (0.4%)	2810 (1.0%)	97 (0.4%)
Anticoagulant use	15 212 (5.3%)	848 (2.9%)	15 372 (5.2%)	688 (2.8%)
Bleeding disorder	4178 (1.4%)	135 (0.5%)	4189 (1.4%)	124 (0.5%)
Cirrhosis	3200 (1.1%)	306 (1.0%)	3221 (1.1%)	285 (1.1%)
Emphysema	15 262 (5.3%)	1322 (4.5%)	15 406 (5.3%)	1178 (4.7%)
Dementia	8474 (2.9%)	372 (1.3%)	8612 (2.9%)	234 (0.9%)
Disseminated cancer	1468 (0.5%)	42 (0.1%)	1472 (0.5%)	38 (0.2%)
Receiving chemotherapy	661 (0.2%)	30 (0.1%)	673 (0.2%)	18 (0.1%)
Alcoholism	24 939 (8.7%)	2544 (8.6%)	24 278 (8.3%)	3205 (12.9%)
Smoker	64 866 (22.5%)	11 714 (39.7%)	65 914 (22.5%)	10 666 (43.0%)
History of substance abuse	24 709 (8.6%)	8457 (28.6%)	25 324 (8.6%)	7842 (31.6%)
Positive alcohol	63 808 (29.1%)	6103 (25.4%)	61 045 (27.5%)	8866 (42.0%)
Mechanism (penetrating)	30 478 (10.6%)	5149 (17.4%)	31 119 (10.6%)	4508 (18.2%)
Head injury AIS score >2	56 023 (19.4%)	5181 (17.5%)	56 746 (19.4%)	4458 (18.0%)
Chest injury AIS score >2	45 528 (15.8%)	5449 (18.4%)	46 888 (16.0%)	4089 (16.5%)
Abdominal injury AIS score >2	13 973 (4.8%)	2093 (7.1%)	14 525 (5.0%)	1541 (6.2%)
Extremity injury AIS score >2	41 762 (14.5%)	4806 (16.3%)	43 017 (14.7%)	3551 (14.3%)
Major surgery	106 408 (36.9%)	12 520 (42.4%)	108 701 (37.1%)	10 227 (41.2%)
Immediate major surgery	32 560 (11.3%)	4223 (14.3%)	33 438 (11.4%)	3345 (13.5%)
Age, mean	45.79	41.26	45.72	41.19
BMI, mean	27.2	26.67	27.19	26.76
Total GCS, mean	13.7	13.5	13.7	13.5
Injury Severity Score, mean	10.3	11.1	10.4	10.6

AIS, Abbreviated Injury Scale; BMI, body mass index; GCS, Glasgow Coma Scale.

were reported with the UDS results. At the time of analysis, it was the latest year available.

Surgery is classified as major according to a database of International Classification of Diseases-10th Revision procedure codes provided by the Agency for Healthcare Research and Quality.⁹ Major surgery is defined as 'procedures that are considered operating room procedures'. Procedures are classified as immediate based on the emergency department disposition. If a patient is transferred directly from the emergency department to the operating room, the procedure is considered immediate. Cardiac and non-cardiac comorbidities are analyzed as individual categorical variables rather than an index. Patients were classified as positive for amphetamine or cocaine independently of UDS results for other drugs, including patients positive for both amphetamine and cocaine.

Analysis was performed for all drug-tested patients. Baseline characteristics are compared for tested versus non-tested patients, and for patients who are positive versus negative for amphetamine and cocaine. Univariate analysis was performed for all drugs included with the UDS results versus all complications reported

in the registry. Further univariate analysis of stimulant drugs was performed for the subsets including all drug-tested patients, those who underwent surgery, and those who underwent immediate emergency surgery. For the stimulant drug analysis, predictor variables were UDS-positive results for methamphetamine and cocaine. Outcome variables were mortality, MI, and stroke. X^2 was used to test for significance for categorical variables.

Multivariate analysis was performed using a multivariate logistic regression with predictor variables of positive amphetamine and cocaine UDS results, age, sex, race, cardiac and non-cardiac comorbidities reported by the NTDB, Injury Severity Score (ISS), mechanism of trauma, trauma regions of head, chest, abdomen, and extremity, surgery, and emergency surgery.

Statistics were performed with the SPSS V.26.0 statistical package (IBM).

RESULTS

There are 997 970 trauma admissions recorded in the NTDB for 2017. A total of 7813 arrived in the emergency room with

Table 2 Univariate analysis of cocaine and amphetamine vs. MI, stroke, and mortality

Outcome	No				RR	P value	No				RR	P value
	Amp	%	Amp	%			Cocaine	%	Cocaine	%		
All patients	288 154	90.70	29 534	9.30			292 868	92.19	24 820	7.81		
MI	441	0.15	26	0.09	0.58	0.005	450	0.15	17	0.07	0.45	<0.001
Stroke	785	0.27	75	0.25	0.93	0.561	803	0.27	57	0.23	0.84	0.195
Mortality	10 474	3.63	896	3.03	0.83	<0.001	10 682	3.65	688	2.77	0.76	<0.001
All surgical patients	106 408	89.47	12 520	10.53			108 701	91.40	10 227	8.60		
MI	280	0.26	19	0.15	0.58	0.019	287	0.26	12	0.12	0.44	0.005
Stroke	588	0.55	58	0.46	0.84	0.198	601	0.55	45	0.44	0.8	0.138
Mortality	4 464	4.2	411	3.28	0.78	<0.001	4 568	4.2	307	3	0.71	<0.001
Immediate surgery	32 560	88.52	4 223	11.48			33 438	90.91	3 345	9.09		
MI	72	0.22	5	0.12	0.54	0.169	74	0.22	3	0.09	0.41	0.112
Stroke	202	0.62	19	0.45	0.73	0.177	203	0.61	18	0.54	0.89	0.623
Mortality	1 886	5.79	174	4.12	0.71	<0.001	1 938	5.8	122	3.65	0.63	<0.001

Significant protective associations shown in bold.

Amp, amphetamine; MI, myocardial infarction; RR, relative risk.

no signs of life or with signs of life not recorded. These were excluded, leaving 990 157 trauma admissions. A total of 317 688 (32.1%) were tested for drugs. Characteristics of patients who underwent a UDS versus those who did not are presented in online supplemental table 1. Results for the following drugs are recorded: amphetamine, barbiturate, benzodiazepines, cocaine, ecstasy, methadone, opioid, oxycodone, phencyclidine, tricyclic antidepressants, and cannabinoids.

Comparison of baseline characteristics (table 1) shows that users of both methamphetamine and cocaine tend to be younger than non-users. Stimulant drug users have a higher likelihood of being male, having a history of substance abuse, and being a victim of penetrating trauma. In general, rates of chronic disease tend to be lower for stimulant drug users.

Univariate analysis of UDS-positive results for all drugs tested showed a protective association with mortality, except for tricyclic antidepressants and methadone which showed no significant association (online supplemental table 2). Positive UDS results showed varying protective and harmful associations with individual complications. Most had a relative risk close to 1. The drug significantly associated with the greatest number of complications was benzodiazepines, whose only protective association was with mortality.

On univariate analysis cocaine and methamphetamine showed protective association with MI and mortality in all drug-tested patients, and in those who underwent major surgery (table 2). Both drugs also showed protective association with mortality for the group who underwent immediate emergency surgery. There were no harmful associations with cardiovascular complications or mortality for stimulant drugs.

Multivariate analysis (table 3) shows that cocaine has a protective association with mortality. Stimulant drugs have no harmful association with mortality or cardiovascular morbidity. Advancing age, ISS, and various comorbidities show a harmful association with mortality and cardiovascular morbidity. The strongest predictor of cardiovascular morbidity is major surgery.

The addition of blood alcohol level (BAL) to the model did not change the significance of any of the predictor values and had a minimal effect on the ORs. We omitted BAL from the final model because 20.1% of patients screened for drugs were

not screened for alcohol, which would result in the omission of many cases.

DISCUSSION

Use of illicit stimulant drugs has been associated with increased morbidity in both the trauma and perioperative setting.¹⁰ To help manage this risk, UDS is used to identify drug-positive patients. However, a positive UDS does not always correlate with acute drug intoxication. For instance, the UDS for cocaine measures levels of an inactive metabolite, which can remain in the urine up to a week after use, making it an unreliable indicator of acute intoxication.¹¹ This makes risk stratification after a positive UDS challenging. Currently, there are no universal guidelines to evaluate a patient's perioperative risk after a positive UDS, but a result positive for stimulant drugs is widely considered a contraindication for elective general anesthesia. A study conducted by Elkassabany *et al* found that two-thirds of the clinicians surveyed would elect to cancel surgery for these patients, even in the absence of any symptoms of acute intoxication.⁵ Canceling elective surgery can delay care in a vulnerable patient population, resulting in adverse health outcomes. It can also result in financial losses and waste of resources. It is difficult to measure the effect that illicit substance use has on elective perioperative complications, given the reluctance to operate on this population. Trauma patients, however, regularly undergo emergency procedures despite having a positive drug screen, allowing us to examine the impact of substance use on perioperative complications.

Our analysis showed protective or neutral association of stimulant drugs with cardiovascular morbidity or mortality on the overall population, and on the subset who underwent major surgery. This lack of harmful association also applied to the group of patients who underwent immediate surgery. If the effects of the substances measured on a UDS conferred excess cardiovascular risk, one would expect this population to be especially vulnerable because the substances had less time for excretion or metabolism. However, even for the group requiring immediate surgery, stimulant-positive UDS is not associated with increased cardiovascular morbidity or mortality.

Table 3 Multivariate analysis of stimulant drugs vs. mortality and cardiovascular outcomes

Characteristics	Multivariate analysis					
	Mortality		MI		Stroke	
	OR	P value	OR	P value	OR	P value
Amphetamine	0.966	0.405	0.801	0.283	1.023	0.857
Age	1.026	<0.001	1.052	<0.001	1.017	<0.001
Sex, female	0.805	<0.001	0.677	<0.001	1.132	0.106
Race (white vs. non-white)	0.959	0.107	0.915	0.471	1.163	0.054
Angina pectoris	0.521	0.099	1.022	0.983	0	0.995
Myocardial infarction	1.243	0.035	2.441	<0.001	1.114	0.739
Congestive heart failure	1.804	<0.001	1.544	0.008	0.969	0.866
Hypertension	0.774	<0.001	1.372	0.005	1.325	0.002
Diabetes mellitus	1.036	0.321	1.554	<0.001	1.439	<0.001
Peripheral arterial disease	1.777	<0.001	1.365	0.396	1.51	0.261
Stroke	1.142	0.042	1.225	0.34	2.473	<0.001
Chronic renal failure	1.946	<0.001	1.118	0.7	0.945	0.851
Anticoagulant use	1.209	<0.001	1.673	<0.001	1.365	0.015
Bleeding disorder	1.118	0.138	0.91	0.752	1.601	0.025
Cirrhosis	3.207	<0.001	1.031	0.934	0.847	0.594
Emphysema	1.593	<0.001	1.408	0.016	1.029	0.843
Dementia	1.016	0.788	0.64	0.064	0.599	0.036
Disseminated cancer	2.156	<0.001	0.973	0.953	1.087	0.833
Receiving chemotherapy	1.682	0.001	1.388	0.582	1.783	0.217
Alcoholism	0.991	0.829	1.09	0.611	1.062	0.628
Smoker	0.512	<0.001	1.632	<0.001	0.936	0.478
Chronic substance abuse	0.767	<0.001	0.865	0.487	0.915	0.495
Injury Severity Score	1.127	<0.001	1.039	<0.001	1.048	<0.001
Mechanism (penetrating vs. blunt)	3.853	<0.001	1.113	0.631	1.084	0.545
Head injury AIS score >2	2.987	<0.001	1.221	0.111	2.594	<0.001
Chest injury AIS score >2	0.782	<0.001	1.366	0.011	1.377	<0.001
Abdominal injury AIS score >2	0.633	<0.001	1.575	0.01	1.072	0.551
Extremity injury AIS score >2	0.687	<0.001	1.144	0.263	0.955	0.595
Had major surgery during admission	0.697	<0.001	2.901	<0.001	4.075	<0.001
Had immediate major surgery	1.360	<0.001	1.124	0.41	1.307	0.002
Cocaine	0.902	0.028	0.627	0.065	0.908	0.502
Age	1.026	<0.001	1.052	<0.001	1.017	<0.001
Sex, female	0.804	<0.001	0.675	<0.001	1.13	0.11
Race (white vs. non-white)	0.963	0.144	0.937	0.598	1.166	0.05
Angina pectoris	0.52	0.098	1.015	0.988	0	0.995
Myocardial infarction	1.243	0.035	2.434	<0.001	1.114	0.74
Congestive heart failure	1.804	<0.001	1.544	0.008	0.968	0.864
Hypertension	0.774	<0.001	1.375	0.005	1.324	0.002
Diabetes mellitus	1.035	0.327	1.554	<0.001	1.439	<0.001
Peripheral arterial disease	1.777	<0.001	1.364	0.398	1.509	0.262
Stroke	1.142	0.041	1.229	0.332	2.473	<0.001
Chronic renal failure	1.942	<0.001	1.113	0.713	0.944	0.847
Anticoagulant use	1.209	<0.001	1.675	<0.001	1.364	0.015
Bleeding disorder	1.116	0.144	0.908	0.744	1.596	0.026
Cirrhosis	3.21	<0.001	1.031	0.934	0.847	0.594
Emphysema	1.594	<0.001	1.409	0.016	1.029	0.843
Dementia	1.015	0.801	0.639	0.064	0.598	0.036
Disseminated cancer	2.154	<0.001	0.974	0.954	1.085	0.836
Receiving chemotherapy	1.68	0.002	1.381	0.588	1.779	0.219
Alcoholism	0.992	0.847	1.095	0.593	1.061	0.633
Smoker	0.513	<0.001	1.644	<0.001	0.94	0.508
Chronic substance abuse	0.773	<0.001	0.892	0.583	0.933	0.595

Continued

Table 3 Continued

Characteristics	Multivariate analysis					
	Mortality		MI		Stroke	
	OR	P value	OR	P value	OR	P value
Injury Severity Score	1.127	<0.001	1.039	<0.001	1.048	<0.001
Mechanism (penetrating vs. blunt)	3.86	<0.001	1.123	0.603	1.088	0.526
Head injury AIS score >2	2.988	<0.001	1.219	0.115	2.593	<0.001
Chest injury AIS score >2	0.782	<0.001	1.364	0.012	1.377	<0.001
Abdominal injury AIS score >2	0.632	<0.001	1.572	0.011	1.072	0.549
Extremity injury AIS score >2	0.687	<0.001	1.142	0.269	0.955	0.599
Had major surgery during admission	0.698	<0.001	2.902	<0.001	4.077	<0.001
Had immediate major surgery	1.359	<0.001	1.121	0.423	1.306	0.002

AIS, Abbreviated Injury Scale; MI, myocardial infarction.

While these findings are consistent with our previous study,⁸ they still seem counterintuitive given the evidence of cardiotoxicity with stimulant use.^{3,4} The protective effects observed with stimulant use may be a result of unknown confounding variables; however, it is also possible that these protective effects are based on a true pharmacologic effect. Evidence of protective effects with stimulant use in trauma patients has been observed in previous studies. Ryb and Cooper showed that patients who had a positive UDS for cocaine had decreased rates of cardiovascular complications when undergoing surgery during the first day of admission¹² and Cheng *et al* showed amphetamine use to be associated with lower rates of mortality.¹ There are also studies that show no significant difference in mortality or cardiovascular complications in trauma patients who test positive for stimulants.^{13–16} This is further validated by recent studies examining the incidence of hemodynamic events under anesthesia to be similar between patients who screen positive for stimulants and patients who have a negative UDS, with the rates of vasopressor use during surgery similar between both groups.^{17–19}

It can be difficult to separate the effects of acute and chronic stimulant use. While we do not know the duration of substance use for each patient, 28.6% and 31.6% of the amphetamine and cocaine group, respectively, in our population had a documented history of chronic substance use. Chronic substance abuse could potentially have a harmful or protective effect. On one hand, chronic stimulant use can result in dilated cardiomyopathy and an increased risk of acute coronary syndrome.^{3,4} However, an alternative hypothesis is that chronic users of methamphetamine and cocaine may have undergone a selection process that leaves survivors less vulnerable to catecholamine-induced cardiovascular stress. Having already survived a stimulant stressor, this population may be relatively resistant to further cardiovascular compromise. Most of our UDS-positive patients did not have a diagnosis of chronic substance abuse. Since we do not know the timing of substance use based on a positive UDS, it is possible that, in spite of positive results, many of our patients were not suffering from acute or chronic effects of illicit drugs at the time of the trauma and subsequent treatment.

While our study findings do not show an association of stimulant use with increased risk of MI, stroke, and mortality, both stimulants are associated with increased rates of surgical site infections, sepsis, and ventilator-associated pneumonia. Stimulants can cause vasoconstriction, resulting in poor wound healing and more susceptibility to surgical site infections. In addition, chronic stimulant abuse can result in malnutrition which can further increase a patient's susceptibility to infections.²⁰ Other

studies have shown that cocaine can negatively impact the functioning of immune cells and mediators.^{20,21}

Strengths of study

We used a large multicenter database for greater statistical power. We were able to show associations between the use of individual drugs and a wide variety of complications. Our study further investigated the use of stimulant drugs and the interaction with surgical procedures to help assess the influence on perioperative risk.

Limitations of study

Not all patients were evaluated for drug use because facilities did not have standardized criteria for choosing which patients to screen. Inconsistent selection of patients for UDS could potentially introduce bias. The UDS does not provide data about the level of the substance present or the time that the substance was used, thus UDS-positive patients may not have been acutely intoxicated. Our analysis shows lower rates of surgical interventions in positive amphetamine/cocaine groups. It is possible that some higher risk patients were excluded from surgery based on UDS results, which could mask a harmful effect. Terms were included in the multivariate analysis to adjust for surgical treatment, but this could still represent a subtle bias. The multivariate analysis adjusts for baseline imbalance; however, the retrospective nature of the study introduces the possibility of unknown and unmeasured confounding variables, thus it cannot establish cause and effect.

CONCLUSION

Our study showed that a UDS positive for cocaine or methamphetamine is not an adverse risk factor for death, stroke, or MI in trauma. This was true for the overall trauma population as well as for the patients who underwent surgery.

This lack of harmful association calls into question the perceived perioperative risk of these drugs and the policy of canceling surgery based on a positive UDS. The results of UDS for illicit drugs show a variety of harmful and protective associations with mortality and complications in trauma patients. It is likely to represent a complex interaction of patient selection and pharmacologic effects. In the setting of a positive UDS, evaluation for signs of acute intoxication, cardiovascular compromise and hemodynamic instability should help guide evaluation of cardiovascular risk and the timing of surgery.

Contributors SS and JC gathered, analyzed and interpreted the patient data regarding the treatment groups and outcomes. CF participated in analyzing the data and interpreting its clinical significance. All authors read and approved the final article.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Our data are from the National Trauma Data Bank. The data were purchased by our institution from the American College of Surgeons

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Cheng V, Inaba K, Johnson M, Byerly S, Jiang Y, Matsushima K, Haltmeier T, Benjamin E, Lam L, Demetriades D. The impact of pre-injury controlled substance use on clinical outcomes after trauma. *J Trauma Acute Care Surg* 2016;81:913–20.
- Beaulieu P. Anesthetic implications of recreational drug use. *Can J Anaesth* 2017;64:1236–64.
- Kim ST, Park T. Acute and chronic effects of cocaine on cardiovascular health. *Int J Mol Sci* 2019;20:584. [Epub ahead of print: 29 Jan 2019].
- Schwarzbach V, Lenk K, Laufs U. Methamphetamine-related cardiovascular diseases. *ESC Heart Fail* 2020;7:407–14.
- Elkassabany N, Speck RM, Oslin D, Hawn M, Chaichana K, Sum-Ping J, Sepulveda J, Whitley M, Sakawi Y. Preoperative screening and case cancellation in cocaine-abusing veterans scheduled for elective surgery. *Anesthesiol Res Pract* 2013;2013:1–7.
- London JA, Battistella FD. Testing for substance use in trauma patients: are we doing enough? *Arch Surg* 2007;142:633–8.
- Soderstrom CA, Dischinger PC, Kerns TJ, Kufera JA, Mitchell KA, Scalea TM. Epidemic increases in cocaine and opiate use by trauma center patients: documentation with a large clinical toxicology database. *J Trauma* 2001;51:557–64.
- Culhane JT, Freeman CA. The effect of illegal drug screening results and chronic drug use on perioperative complications in trauma. *J Emerg Trauma Shock* 2020;13:279–85.
- Procedure Classes Refined for ICD-10-PCS. *Healthcare cost and utilization project (HCUP)*. Rockville, MD: Agency for Healthcare Research and Quality, 2020.
- Vadivelu N, Mitra S, Kaye AD, Urman RD. Perioperative analgesia and challenges in the drug-addicted and drug-dependent patient. *Best Pract Res Clin Anaesthesiol* 2014;28:91–101.
- Saggese NP, Chang C, Cardo VA. Perioperative management for the cocaine-positive patient undergoing elective surgery under general anesthesia. *J Oral Maxillofac Surg* 2019;77:894–5.
- Ryb GE, Cooper C. Outcomes of cocaine-positive trauma patients undergoing surgery on the first day after admission. *J Trauma* 2008;65:809–12.
- Fujii Q, McCague A. Recreational drugs and outcomes in trauma patients. *J Emerg Trauma Shock* 2019;12:98–100.
- Hadjizacharia P, Green DJ, Plurad D, Chan LS, Inaba K, Shulman I, Demetriades D. Methamphetamines in trauma: effect on injury patterns and outcome. *J Trauma* 2009;66:895–8.
- Hadjizacharia P, Green DJ, Plurad D, Chan LS, Law J, Inaba K, Demetriades D. Cocaine use in trauma: effect on injuries and outcomes. *J Trauma* 2009;66:491–4.
- Yeung JT, Williams J, Bowling WM. Effect of cocaine use on outcomes in traumatic brain injury. *J Emerg Trauma Shock* 2013;6:189–94.
- Moon TS, Gonzales MX, Sun JJ, Kim A, Fox PE, Minhajuddin AT, Pak TJ, Ogunnaike B. Recent cocaine use and the incidence of hemodynamic events during general anesthesia: a retrospective cohort study. *J Clin Anesth* 2019;55:146–50.
- Edwards AM, Johnson EG, Bernard AC. Intraoperative vasopressor use during emergency surgery on injured Meth users. *Trauma Surg Acute Care Open* 2020;5:e000553.
- Hill GE, Ogunnaike BO, Johnson ER. General anaesthesia for the cocaine abusing patient. is it safe? *Br J Anaesth* 2006;97:654–7.
- Brownlow HA, Pappachan J. Pathophysiology of cocaine abuse. *Eur J Anaesthesiol* 2002;19:395.
- Baldwin GC, Tashkin DP, Buckley DM, Park AN, Dubinett SM, Roth MD. Marijuana and cocaine impair alveolar macrophage function and cytokine production. *Am J Respir Crit Care Med* 1997;156:1606–13.

Characteristics	Drug Tested	Not Drug Tested
Sex, Female	105491(33.2%)	290529(43.2%)
Race (White vs Non-white)	217508(68.5%)	506758(75.4%)
Hispanic Ethnicity	18075(30.8%)	30961(30.5%)
Angina Pectoris	240(0.1%)	810(0.1%)
Myocardial Infarction	2240(0.7%)	6485(1.0%)
Congestive Heart Failure	8398(2.6%)	28650(4.3%)
Hypertension	82359(25.9%)	227047(33.8%)
Diabetes Mellitus	32835(19.3%)	89112(13.3%)
Peripheral Arterial Disease	1102(0.3%)	3876(0.6%)
Stroke	6145(1.9%)	19008(2.8%)
Chronic Renal Failure	2907(0.9%)	12373(1.8%)
Anticoagulant Use	16060(5.1%)	57286(8.5%)
Bleeding Disorder	4313(1.4%)	10944(1.6%)
Cirrhosis	3506(1.1%)	6115(0.9%)
Emphysema	16584(5.2%)	44525(6.6%)
Dementia	8846(2.8%)	42909(6.4%)
Disseminated Cancer	1510(0.5%)	3982(0.6%)
Receiving Chemotherapy	691(0.2%)	2805(0.4%)
Alcoholism	27483(8.7%)	24968(3.7%)
Smoker	76580(24.1%)	100919(15.0%)
History of Substance Abuse	33166(10.4%)	17865(2.7%)
Positive Alcohol on Presentation	69911(28.7%)	45819(26.1%)
Mechanism (Penetrating vs Blunt)	35627(11.2%)	52061(7.7%)
Head Injury AIS>2	61204(19.3%)	92352(13.7%)
Chest Injury AIS>2	50977(16.0%)	67289(10.0%)
Abdominal Injury AIS>2	16066(5.1%)	19381(2.9%)
Extremity Injury AIS>2	46568(14.7%)	128879(19.2%)
Major Surgery During Admission	118928(37.4%)	270610(40.2%)
Had Immediate Major Surgery	36783(11.6%)	71317(10.6%)
Age, mean	45.4	48.8
BMI, mean	27.2	26.6
Total GCS mean	13.7	14.5
Injury Severity Score mean	10.4	8.3

Supplemental Table 1. Baseline Characteristics of Patients Tested for Drugs Versus Those Not Tested

Complication	All				RR		Cocaine			
	All	%	Amp	%	RR	p	Cocaine	%	RR	p
CLABSI	190	0.06	23	0.08	1.34	0.18	18	0.07	1.23	0.39
Deep SSI	427	0.13	62	0.21	1.66	<0.01	52	0.21	1.64	<0.01
DVT	2283	0.72	280	0.95	1.36	<0.01	199	0.8	1.13	0.11
EtOH										
Withdrawal	2770	0.87	206	0.7	0.78	<0.01	248	1	1.16	0.03
Cardiac Arrest	2009	0.63	173	0.59	0.92	0.29	147	0.59	0.93	0.41
CAUTI	881	0.28	115	0.39	1.46	<0.01	92	0.37	1.38	<0.01
Embolism	1063	0.33	103	0.35	1.05	0.66	99	0.4	1.21	0.07
Extremity CS	322	0.1	43	0.15	1.5	0.01	26	0.1	1.04	0.86
Intubation	3538	1.11	361	1.22	1.11	0.06	294	1.18	1.07	0.27
Kidney	1764	0.56	195	0.66	1.21	0.01	155	0.62	1.14	0.13
MI	467	0.15	26	0.09	0.58	0.01	17	0.07	0.45	<0.01
Organ Space										
SSI	298	0.09	43	0.15	1.65	<0.01	38	0.15	1.72	<0.01
Osteomyelitis	64	0.02	8	0.03	1.39	0.38	7	0.03	1.45	0.35
Other	20267	6.38	4138	14.01	2.5	<0.01	3606	14.53	2.55	<0.01
Respiratory	1356	0.43	125	0.42	0.99	0.92	117	0.47	1.11	0.26
Return to OR	1630	0.51	170	0.58	1.14	0.11	146	0.59	1.16	0.08
Sepsis	1105	0.35	131	0.44	1.31	<0.01	110	0.44	1.3	0.01
Stroke	860	0.27	75	0.25	0.93	0.56	57	0.23	0.84	0.20
Superficial SSI	350	0.11	43	0.15	1.37	0.05	44	0.18	1.7	<0.01
Pressure Ulcer	1278	0.4	114	0.39	0.96	0.64	108	0.44	1.09	0.39
Unplanned										
RICU	4291	1.35	385	1.3	0.96	0.46	358	1.44	1.07	0.19
VAP	2206	0.69	286	0.97	1.45	<0.01	245	0.99	1.47	<0.01
Mortality	11370	3.58	896	3.03	0.83	<0.01	688	2.77	0.76	<0.01

Complication	Benzo				Barb		Ecsy					
	Benzo	%	RR	p	Barb	%	RR	p	Ecsy	%	RR	p
CLABSI	12	0.05	0.89	0.70	5	0.08	1.3	0.56	3	0.08	1.31	0.64
Deep SSI	52	0.23	1.83	<0.01	12	0.19	1.39	0.26	9	0.23	1.76	0.09
DVT	195	0.87	1.23	0.01	22	0.34	0.47	<0.01	8	0.21	0.29	<0.01
EtOH												
Withdrawal	277	1.24	1.47	<0.01	56	0.87	0.99	0.97	14	0.36	0.41	<0.01
Cardiac Arrest	140	0.63	0.99	0.90	25	0.39	0.61	0.01	6	0.16	0.24	<0.01
CAUTI	100	0.45	1.69	<0.01	45	0.7	2.59	<0.01	37	0.96	3.58	<0.01
Embolism	100	0.45	1.37	<0.01	21	0.33	0.97	0.89	12	0.31	0.93	0.81
Extremity CS	35	0.16	1.61	0.01	4	0.06	0.61	0.31	1	0.03	0.25	0.14
Intubation	302	1.35	1.23	<0.01	84	1.3	1.17	0.15	45	1.17	1.05	0.74
Kidney	150	0.67	1.23	0.02	29	0.45	0.81	0.25	20	0.52	0.94	0.77
MI	24	0.11	0.71	0.11	7	0.11	0.73	0.41	0	0	0	0.02

Organ Space												
SSI	30	0.13	1.48	0.04	6	0.09	0.99	0.98	4	0.1	1.11	0.84
Osteomyelitis	7	0.03	1.62	0.22	0	0	0	0.25	0	0	0	0.38
Other	3793	16.95	3.04	<0.01	2726	42.2	7.49	<0.01	2574	66.93	11.87	<0.01
Respiratory	156	0.7	1.72	<0.01	10	0.15	0.36	<0.01	2	0.05	0.12	<0.01
Return to OR	157	0.7	1.41	<0.01	19	0.29	0.57	0.01	13	0.34	0.66	0.13
Sepsis	112	0.5	1.49	<0.01	26	0.4	1.16	0.45	15	0.39	1.12	0.66
Stroke	62	0.28	1.03	0.85	15	0.23	0.86	0.55	10	0.26	0.96	0.90
Superficial SSI	40	0.18	1.7	<0.01	4	0.06	0.56	0.24	2	0.05	0.47	0.27
Pressure Ulcer	97	0.43	1.08	0.44	11	0.17	0.42	<0.01	3	0.08	0.19	<0.01
Unplanned												
RICU	345	1.54	1.15	0.01	95	1.47	1.09	0.40	72	1.87	1.39	0.01
VAP	276	1.23	1.89	<0.01	63	0.98	1.42	0.01	53	1.38	2.01	<0.01
Mortality	724	3.24	0.9	<0.01	194	3	0.84	0.01	98	2.55	0.71	<0.01
	Mdn	%	RR	p	Opioid	%	RR	p	Oxy	%	RR	p
CLABSI	4	0.16	2.76	0.04	8	0.04	0.58	0.13	2	0.06	0.95	0.95
Deep SSI	3	0.12	0.91	0.87	34	0.15	1.14	0.47	4	0.11	0.85	0.74
DVT	12	0.49	0.68	0.17	117	0.52	0.71	<0.01	13	0.37	0.51	0.01
EtOH												
Withdrawal	29	1.18	1.36	0.10	193	0.86	0.99	0.85	34	0.97	1.12	0.53
Cardiac Arrest	13	0.53	0.83	0.52	93	0.41	0.64	<0.01	10	0.29	0.45	0.01
CAUTI	12	0.49	1.77	0.05	49	0.22	0.78	0.08	13	0.37	1.34	0.29
Embolism	9	0.37	1.09	0.79	77	0.34	1.03	0.82	6	0.17	0.51	0.09
Extremity CS	6	0.24	2.43	0.03	35	0.16	1.61	0.01	4	0.11	1.13	0.81
Intubation	38	1.55	1.39	0.04	219	0.98	0.87	0.04	46	1.31	1.18	0.26
Kidney	15	0.61	1.1	0.71	108	0.48	0.86	0.12	12	0.34	0.61	0.09
MI	4	0.16	1.11	0.84	25	0.11	0.74	0.15	3	0.09	0.58	0.34
Organ Space												
SSI	1	0.04	0.43	0.39	20	0.09	0.95	0.81	1	0.03	0.3	0.21
Osteomyelitis	0	0	0	0.48	5	0.02	1.12	0.81	0	0	0	0.40
Other	103	4.19	0.65	<0.01	1289	5.75	0.89	<0.01	166	4.74	0.74	<0.01
Respiratory	10	0.41	0.95	0.88	82	0.37	0.85	0.14	6	0.17	0.4	0.02
Return to OR	5	0.2	0.39	0.03	96	0.43	0.82	0.06	24	0.69	1.34	0.15
Sepsis	13	0.53	1.53	0.13	74	0.33	0.94	0.64	6	0.17	0.49	0.08
Stroke	8	0.33	1.2	0.60	43	0.19	0.69	0.02	8	0.23	0.84	0.63
Superficial SSI	3	0.12	1.11	0.86	24	0.11	0.97	0.88	5	0.14	1.3	0.56
Pressure Ulcer	9	0.37	0.91	0.78	49	0.22	0.52	<0.01	8	0.23	0.57	0.10
Unplanned												
RICU	30	1.22	0.9	0.57	286	1.27	0.94	0.31	51	1.46	1.08	0.59
VAP	20	0.81	1.17	0.48	113	0.5	0.71	<0.01	21	0.6	0.86	0.50
Mortality	84	3.42	0.95	0.66	425	1.89	0.51	<0.01	81	2.31	0.64	<0.01
	PCP	%	RR	p	Tricyc	%	RR	p	Cnb	%	RR	p

CLABSI	4	0.14	2.4	0.07	2	0.13	2.11	0.28	35	0.06	1.04	0.84
Deep SSI	4	0.14	1.06	0.91	2	0.13	0.93	0.92	97	0.17	1.35	0.01
DVT	17	0.6	0.84	0.47	10	0.63	0.87	0.66	474	0.84	1.21	<0.01
EtOH												
Withdrawal	22	0.78	0.89	0.60	27	1.69	1.95	<0.01	535	0.94	1.1	0.04
Cardiac Arrest	19	0.67	1.07	0.78	6	0.38	0.59	0.20	379	0.67	1.07	0.23
CAUTI	12	0.43	1.54	0.13	9	0.56	2.04	0.03	143	0.25	0.89	0.21
Embolism	13	0.46	1.38	0.24	3	0.19	0.56	0.31	185	0.33	0.97	0.70
Extremity CS	2	0.07	0.7	0.61	3	0.19	1.86	0.28	86	0.15	1.68	<0.01
Intubation	43	1.53	1.37	0.04	27	1.69	1.52	0.03	607	1.07	0.95	0.28
Kidney	21	0.75	1.35	0.17	14	0.88	1.58	0.08	263	0.46	0.81	<0.01
MI	5	0.18	1.21	0.67	3	0.19	1.28	0.67	40	0.07	0.43	<0.01
Organ Space												
SSI	2	0.07	0.75	0.69	0	0	0	0.22	74	0.13	1.52	<0.01
Osteomyelitis	0	0	0	0.45	0	0	0	0.57	16	0.03	1.53	0.14
Other	90	3.19	0.5	<0.01	66	4.13	0.65	<0.01	2814	4.96	0.74	<0.01
Respiratory	14	0.5	1.17	0.57	9	0.56	1.32	0.40	289	0.51	1.25	<0.01
Return to OR	18	0.64	1.25	0.35	9	0.56	1.1	0.78	411	0.72	1.55	<0.01
Sepsis	16	0.57	1.64	0.05	3	0.19	0.54	0.28	207	0.37	1.06	0.44
Stroke	8	0.28	1.05	0.89	6	0.38	1.39	0.42	113	0.2	0.7	<0.01
Superficial SSI	6	0.21	1.95	0.10	0	0	0	0.18	87	0.15	1.52	<0.01
Pressure Ulcer	13	0.46	1.15	0.62	6	0.38	0.93	0.87	242	0.43	1.07	0.31
Unplanned												
RICU	30	1.06	0.79	0.19	19	1.19	0.88	0.58	640	1.13	0.81	<0.01
VAP	26	0.92	1.33	0.14	15	0.94	1.36	0.24	500	0.88	1.35	<0.01
Mortality	79	2.8	0.78	0.03	61	3.82	1.07	0.60	1635	2.88	0.77	<0.01

Supplemental Table 2. Univariate Analysis of Illicit Drug UDS with Complications. (Significant protective and harmful associations shown in blue and red, respectively). Amp: Amphetamine, Barb: Barbiturate, Ecsy: Ecstasy, Benzo: Benzodiazepine, Mdn: Methadone, Oxy: Oxycodone, PCP: Phencyclidine, Tricyc: Tricyclic Antidepressant, Cnb: Cannabinoid

CLABSI: Central Line Associated Bloodstream Infection, SSI: Surgical Site Infection, DVT: Deep Venous Thrombosis, CAUTI: Catheter Associated Urinary Tract Infection, CS: Compartment Syndrome, RICU: Return to Intensive Care Unit, VAP: Ventilator Associated Pneumonia