

# The national trauma triage protocol: how EMS perspective can inform the guideline revision

Peter E Fischer,<sup>1</sup> Mark L Gestring,<sup>2</sup> Scott G Sagraves,<sup>3</sup> Holly N Michaels,<sup>4</sup> Bhavin Patel,<sup>4</sup> Jimm Dodd,<sup>4</sup> Eric M Campion,<sup>5</sup> Wayne E VanderKolk,<sup>6</sup> Eileen M Bulger<sup>7</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/tsaco-2021-000879>).

<sup>1</sup>Department of Surgery, The University of Tennessee Health Science Center, Memphis, Tennessee, USA

<sup>2</sup>Department of Surgery, University of Rochester, Rochester, New York, USA

<sup>3</sup>Department of Surgery, Baylor Scott and White Central Texas, Temple, Texas, USA

<sup>4</sup>Committee on Trauma, American College of Surgeons, Chicago, Illinois, USA

<sup>5</sup>Department of Surgery, Denver Health, Denver, Colorado, USA

<sup>6</sup>Department of Surgery, West Michigan Surgical Specialists, Grand Rapids, Michigan, USA

<sup>7</sup>Department of Surgery, University of Washington, Seattle, Washington, USA

## Correspondence to

Dr Peter E Fischer; [pfischer@uthsc.edu](mailto:pfischer@uthsc.edu)

The 80th Annual Meeting of the American Association for the Surgery of Trauma & Clinical Congress of Acute Care Surgery

Received 27 December 2021  
Accepted 31 December 2021

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Fischer PE, Gestring ML, Sagraves SG, et al. *Trauma Surg Acute Care Open* 2022;**7**:e000879.

## ABSTRACT

**Objectives** The Field Triage Guidelines (FTG) support emergency medical service (EMS) decisions regarding the most appropriate transport destination for injured patients. While the components of the algorithm are largely evidenced-based, the stepwise approach was developed with limited input from EMS providers. FTG are only useful if they can easily be applied by the field practitioner. We sought to gather end-user input on the current guidelines from a broad group of EMS stakeholders to inform the next revision of the FTG.

**Methods** An expert panel composed an end-user feedback tool. Data collected included: demographics, EMS agency type, geographic area of respondents, use of the current FTG, perceived utility, and importance of each step in the algorithm (1: physiologic, 2: anatomic, 3 mechanistic, 4: special populations). The American College of Surgeons Committee on Trauma (ACS COT), in partnership with several key organizations, distributed the tool to reach as many providers as possible.

**Results** 3958 responses were received (82% paramedics/emergency medical technicians, 9% physicians, 9% other). 94% responded directly to scene emergency calls and 4% were aeromedical providers. Steps 2 and 3 were used in 95% of local protocols, steps 1 and 4 in 90%. Step 3 was used equally in protocols across all demographics; however, step 1 was used significantly more in the air medical services than ground EMS (96% vs 88%,  $p < 0.05$ ). Geographic variation was demonstrated in FTG use based on the distance to a trauma center, but step 3 (not step 1) drove the majority of the decisions. This point was reinforced in the qualitative data with the comment, "I see the wreck before I see the patient."

**Conclusion** The FTG are widely used by EMS in the USA. The stepwise approach is useful; however, mechanism (not physiological criteria) drives most of the decisions and is evaluated first. Revision of the FTG should consider the experience of the end-users.

**Level of evidence** V.

## INTRODUCTION

The Guidelines for Field Triage of Injured Patients (FTG) were originally developed to assist emergency medical service (EMS) providers identify injured patients at the scene who would benefit from trauma center care and support EMS transport destination decisions. The FTG were originally developed in 1976 by the American College of Surgeons, with periodic revisions every 5–10 years.<sup>1</sup> The most recent revisions of the FTG were led by

the US Centers for Disease Control and completed in 2011.<sup>2</sup>

The FTG were designed to be used in a stepwise fashion such that physiological criteria were evaluated first (step 1), followed by anatomic criteria (step 2), then mechanism of injury criteria (step 3) and lastly, special considerations related to specific patient populations (step 4). While the components of this algorithm were largely evidence-based or driven by expert consensus when evidence was not available, these guidelines were developed with limited input from the EMS providers who use them in the field. With this in mind, we actively sought input regarding the current guidelines from a broad group of EMS stakeholders by distributing an end-user feedback tool throughout the EMS community to gather input for future revisions of the FTG.

## METHODS

An expert panel composed of nine members of the EMS subcommittee of the American College of Surgeons Committee on Trauma drafted a feedback tool consisting of 40 questions (online supplemental appendix A). The tool was separated into sections examining responder characteristics and FTG use specifically by step. The demographic section covered the respondent's role in EMS, the type of service where the respondent was currently working, and general geographic data based on the transport time to a trauma center. General questions were next and examined the perceived use of the current guidelines and the role of EMS judgment. Each step of the 2011 algorithm was then assessed by the next set of questions probing ease of use, importance in triage, and suggestions for improvement. For ease of analysis, questions, when possible, were limited to quantitative responses, however qualitative free-text answers were also allowed.

The feedback tool was then piloted in a small group of EMS providers to assess question understanding and refine any areas which caused confusion. Once the tool was complete, the goal was to widely distribute the tool to a diverse array of EMS providers from across the USA. This was accomplished by partnering with major national EMS organizations to widely distribute the feedback tool to their members via print, email, and social media (online supplemental appendix B). The tool was open for responses for approximately a 6-week period in the fall of 2020 on the Qualtrics platform and allowed respondents to answer the questions using smart phone or computer. Respondents were

**Table 1** Primary role of respondent in EMS

Primary role at EMS	%	Count
Patient care provider	58	2231
Administrator/Manager	15	572
First-line supervisor	14	525
Educator	7	275
Other	5	181
Preceptor	1	29

EMS, emergency medical service.

not stopped from answering the questions twice and partial completion of the tool was permitted. Data was collected in the Qualtrics platform and analyzed using Qualtrics Stats iQ software and PivotTables. Free-text answers were individually reviewed by the expert panel and organized by theme.

## RESULTS

A total of 3958 responses to the end-user feedback tool were submitted. Given that the tool was distributed widely across many organizations, a true response rate could not be calculated. The demographics of the respondents are shown in [table 1](#). Most of the respondents were paramedics/emergency medical technicians and 94% responded to scene calls.

All steps of the current guideline (step 1: physiological parameters; step 2: anatomic factors; step 3: mechanisms of injury; step 4: special considerations) were used in local protocols (step 1 89%, step 2 94%, step 3 93%, and step 4 90%). Interestingly, step 1 was used more often in aeromedical service protocols than in ground services. The role of EMS judgment when applied to the FTG was also examined. Most respondents (62%) replied that their judgment overrode the FTG only 0%–20% of the time. When judgment did override, most of the time (52%) a patient was transported to a trauma center when the patient did not meet other criteria. Only 8% of the time did the respondents transport patients to a non-trauma center who met criteria to be triaged to a trauma center based on their judgment.

Eighty-eight per cent of the respondents felt the stepwise approach to trauma triage was useful. The importance of each step as judged by the respondent is shown in [table 2](#). The order of the steps was also examined. Steps 1 and 3 were preferentially ranked number 1 when determining a patient transport destination. Steps 2 and 3 were ranked evenly at number 3 ([table 3](#)). When asked which step drives the *majority* of the decisions to triage a patient to a trauma center, the respondents chose step 3 which represents the mechanism of injury.

The respondents practiced in systems with variable access to trauma centers. Eighty-four per cent of the respondents said a major trauma patient in their area would be taken to the closest level 1 or 2 center, while only 6% said the closest hospital (non-trauma center). The average time to reach a level 1 or 2 trauma

**Table 2** Importance of each step of the criteria for appropriate triage as judged by the respondent

Triage importance	Step 1 (%)	Step 2 (%)	Step 3 (%)	Step 4 (%)
Extremely	45	40	26	23
Very	38	48	41	41
Moderately	14	10	27	28
Slightly	2	1	5	7
Not	1	1	1	1

center by ground was 29 min (range 0.1– 800). The average time to reach any level trauma center was lower at 19 min (range 0.1–500). When asked how often a major trauma patient would be transported by air, over 70% replied only 0%–25% of the time. When stratifying the responses by time to a trauma center, the patient was more likely to be triaged to a non-trauma center and air transport used more often the further away EMS was from a trauma center ([table 4](#)). Step 3 drove the majority of the triage decisions regardless of transport time to the trauma center.

Almost all questions on the end-user feedback tool provided the opportunity to place free-text responses, and over 6000 free-text answers were received. These responses were each individually evaluated and grouped into categories as best as possible. Data continued to suggest that while users felt the stepwise approach was easy to understand, the flow and usefulness of the tool in the field did not match the order of the steps in the guideline. Users provided feedback that step 3 was often used first in field triage. Furthermore, suggestions included simplifying step 1 by removing Glasgow Coma Scale (GCS) (as it is difficult to remember in the field) and adding age-specific physiological criteria. Multiple comments requested removing concrete diagnosis (eg, pelvic fractures) from step 2 and favored more ‘suspicion’ since definitive diagnosis cannot be made in the field. Multiple comments also requested clarifying the role of EMS judgment.

## DISCUSSION

We gathered significant amounts of end-user feedback on the 2011 FTG. The feedback was primarily from providers who work in the prehospital setting and use the guidelines regularly. All steps of the current guideline were used in local protocols, although to varying degrees. While the rate of use was high ( $\geq 89\%$ ) for all steps, it certainly was not 100%. State and local EMS Medical Program Directors may make modifications to the guideline for local implementation. This is consistent with previous studies which have described variability in adoptions of the guidelines. Barnett *et al* found the most consistent use of the FTG was the physiological criteria while anatomic criteria were not as consistent.<sup>3</sup> Our results were slightly different demonstrating more use of the anatomic and mechanistic criteria in local protocol than the physiological criteria. When examining state adoption of the FTG 4 years after a major revision, only 17% had fully adopted all criteria.<sup>4</sup> Our study was conducted 8 years after the publication of the 2011 guidelines and is consistent with other studies demonstrating that increased time between survey and introduction will reflect a higher rate of implementation.<sup>5</sup>

An interesting finding was that the step 1 criteria were more likely to be integrated into air-medical protocols than ground services protocols. While the exact reasons for this could not be determined, it is plausible that these providers used the physiological data more often for triage due to inability to assess the severity of the mechanism. Often air-medical teams will rendezvous with ground services who will have the patient already in the ambulance and thus the air medical crews may not see the scene of the injury.

The concept of EMS provider judgment was added to the FTG in the 2006 revision and retained in the 2011 revision, however the parameters to guide decisions made by EMS outside of the algorithm were not described.<sup>2</sup> Sixty-two per cent of our respondents stated that their judgment overrode the FTG criteria <20% of the time. This is contrary to prior research showing that judgment was the most commonly used criterion when EMS providers were asked

**Table 3** The end-user feedback tool asked the respondents to order the steps of the FTGs in terms of importance and use as a provider in the field determining patient transport destination

FTG step	Ranked #1		Ranked #2		Ranked #3		Ranked #4	
	Importance (%)	Use (%)						
1	38	38	31	30	23	24	8	9
2	23	21	43	45	29	29	5	5
3	36	38	22	21	31	31	11	10
4	3	3	4	4	17	16	76	76

FTG, Field Triage Guidelines.

immediately after transporting a patient.<sup>6</sup> Studies regarding the utility of EMS judgment are mixed. Some studies have shown that EMS judgment adds to the sensitivity of triage,<sup>6–8</sup> while others show no benefit.<sup>9,10</sup> Regardless of the accuracy, research shows that the ‘gut feeling’ of the provider is the primary reason for identifying a major trauma victim requiring trauma center triage.<sup>11</sup> Further revisions of the FTGs need to continue to recognize EMS judgment as an important part of the triage process and efforts should be made to help further develop the factors to be considered which may prompt a provider to transport a patient to a trauma center even when they fail to meet other criteria.

Our results show that as driving distance to a trauma center increases, the patient was less likely to be taken to a trauma center. This is despite data showing that major trauma patient transport directly to a trauma center improves survival.<sup>12</sup> There are likely multiple system issues effecting the transport decision. For example, many rural counties have a very limited number of ambulances and cannot allow for an extended transport time to a trauma center at the cost of leaving an area without any EMS support. In these circumstances, an ambulance will often take a major trauma victim to the closest hospital, which is often not a trauma center. These results are not unique. Newgard *et al* found that while the identification of high-risk patients did not differ between urban and rural environments, the transport decisions varied widely.<sup>13,14</sup> Any revision of the FTG must account for trauma center availability and geographical restraints within a system.

Mechanistic criteria are important in the current FTG. However, in the current design of the algorithm, the mechanism is not meant to be considered until after the physiological and anatomic criteria. Thus, if the FTG are used in the correct sequence steps 1 and 2 should drive the majority of triage decisions to a trauma center. However, this is not consistent with our findings of how the FTG are used in the field. Providers consistently ranked step 3 (mechanism) as the primary step driving most of the decisions to take a patient to a trauma center. These findings are consistent with a cognitive reasoning model described.<sup>11</sup> The mechanism criteria are often examined first based on dispatch information and visual cues on scene arrival. This finding was reinforced by one of the comments on the tool. The provider stated, “I see the scene before I see the patient.” The current stepwise approach of the algorithm

is more consistent with the evaluation of the patient in the hospital where vital signs are obtained first as part of the primary survey and are not consistent with the flow of information in the prehospital environment.

There are a number of important limitations to the data presented. Based on our strategy to ensure wide distribution of the tool, we cannot determine the response rate as we do not know the denominator of the number of individuals that received the tool. We also cannot assess the demographics of those who did not submit responses. Furthermore, while the tool was piloted with EMS providers to ensure clarity, it is possible the questions were interpreted differently among respondents. With the large response, we hope that the EMS community was appropriately represented but there is the possibility of sampling error.

In conclusion, the current FTG are widely used across EMS agencies. Each step is well represented in local protocols. However, step 3, not step 1, drove the majority of the decisions to triage a patient to the trauma center likely secondary to the flow of information prehospital professionals receive in the field. Proximity to a trauma center also played a role in how trauma patients were triaged. FTG should be adjusted for field workflow and be adaptable based on specific local system requirements. Overall, end-user input is essential for future guideline revisions.

**Contributors** PF, MG, SS, HM, JD, EC, WV, EB composed the feedback tool and manuscript. BP performed data analysis. PF is the guarantor.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants but was not approved by an Ethics Committee(s) or Institutional Board(s). The survey was conducted to help guide the next revision of the field triage guidelines and was not originally designed to be a research questionnaire.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Data availability statement** Data may be obtained from a third party and are not publicly available.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

**REFERENCES**

- Mackersie RC. History of trauma field triage development and the American College of surgeons criteria. *Prehosp Emerg Care* 2006;10:287–94.
- Sasser SM, Hunt RC, Faul M, Sugarman D, Pearson WS, Dulski T, Wald MM, Jurkovich GJ, Newgard CD, Lerner EB, *et al*. Guidelines for field triage of injured patients: recommendations of the National expert panel on field triage, 2011. *MMWR Recomm Rep* 2012;61:1–20.
- Barnett AS, Wang NE, Sahni R, Hsia RY, Haukoos JS, Barton ED, Holmes JF, Newgard CD. Variation in prehospital use and uptake of the National field triage decision scheme. *Prehosp Emerg Care* 2013;17:135–48.

**Table 4** Non-TC patient transport, use of air medical services, and step of FTGs driving majority of triage decisions stratified by average distance by ground to any level trauma center

Ground time to trauma center (min)	Non-TC transport (%)		Step driving majority of triage decisions
	Non-TC transport (%)	Air transport (%)	
0–30	3	11	3
31–60	22	53	3
61+	59	69	3

FTG, Field Triage Guidelines; TC, trauma center.

- 4 Sasser SM, Ossmann E, Wald MM, Lerner EB, Hunt RC. Adoption of the 2006 field triage decision scheme for injured patients. *West J Emerg Med* 2011;12:275–83.
- 5 Bigham BL, Koprowicz K, Aufderheide TP, Davis DP, Donn S, Powell J, Suffoletto B, Nafziger S, Stouffer J, Idris A, et al. Delayed prehospital implementation of the 2005 American heart association guidelines for cardiopulmonary resuscitation and emergency cardiac care. *Prehosp Emerg Care* 2010;14:355–60.
- 6 Newgard CD, Kampp M, Nelson M, Holmes JF, Zive D, Rea T, Bulger EM, Liao M, Sherck J, Hsia RY, et al. Deciphering the use and predictive value of "emergency medical services provider judgment" in out-of-hospital trauma triage: a multisite, mixed methods assessment. *J Trauma Acute Care Surg* 2012;72:1239–48.
- 7 Engum SA, Mitchell MK, Scherer LR, Gomez G, Jacobson L, Solotkin K, Grosfeld JL. Prehospital triage in the injured pediatric patient. *J Pediatr Surg* 2000;35:82–7.
- 8 Lavoie A, Emond M, Moore L, Camden S, Liberman M. Evaluation of the prehospital index, presence of high-velocity impact and judgment of emergency medical technicians as criteria for trauma triage. *CJEM* 2010;12:111–8.
- 9 Mulholland SA, Gabbe BJ, Cameron P. Is paramedic judgement useful in prehospital trauma triage? *Injury* 2005;36:1298–305.
- 10 Qazi K, Kempf JA, Christopher NC, Gerson LW. Paramedic judgment of the need for trauma team activation for pediatric patients. *Acad Emerg Med* 1998;5:1002–7.
- 11 Newgard CD, Nelson MJ, Kampp M, Saha S, Zive D, Schmidt T, Daya M, Jui J, Wittwer L, Warden C, et al. Out-Of-Hospital decision making and factors influencing the regional distribution of injured patients in a trauma system. *J Trauma* 2011;70:1345–53.
- 12 Haas B, Stukel TA, Gomez D, Zagorski B, De Mestral C, Sharma SV, Rubinfeld GD, Nathens AB. The mortality benefit of direct trauma center transport in a regional trauma system: a population-based analysis. *J Trauma Acute Care Surg* 2012;72:1510–5.
- 13 Newgard CD, Fu R, Bulger E, Hedges JR, Mann NC, Wright DA, Lehrfeld DP, Shields C, Hoskins G, Warden C, et al. Evaluation of rural vs urban trauma patients served by 9-1-1 emergency medical services. *JAMA Surg* 2017;152:11–18.
- 14 Newgard CD, Fu R, Zive D, Rea T, Malveau S, Daya M, Jui J, Griffiths DE, Wittwer L, Sahni R, et al. Prospective validation of the National field triage guidelines for identifying seriously injured persons. *J Am Coll Surg* 2016;222:146–58.

## Appendix A

# Field Triage of Injured Patients Guideline (FTG)– Stakeholder Feedback Tool

---

Start of Block: Introductory Questions- Page 1

### Q1 Demographics

Which of the following best describes your primary role at your main EMS job?

- Patient Care Provider** - A person whose primary role is the provision of emergency medical services to patients.
  - Educator** - A person whose primary role is instructing individuals enrolled in an approved or accredited EMS training course or providing continuing education required for maintenance of licensure.
  - Preceptor** - A person whose primary role is training individuals enrolled in an approved or accredited EMS training course in a clinical setting.
  - Dispatcher/Operator** - A person whose primary role is EMS communications.
  - Administrator/Manager** - A person whose primary role is the management and direction of an organization providing emergency medical services.
  - First-line Supervisor** - A person whose primary role is the direct supervision of individuals providing emergency medical services.
  - Other** - A person whose primary EMS role at their main job is not listed above:  
\_\_\_\_\_
-

Q2 At what level are you currently practicing as an EMS provider?

- Emergency Medical Responder; First Responder
  - Emergency Medical Technician - Basic
  - Emergency Medical Technician - Intermediate
  - Paramedic
  - Advanced Emergency Medical Technician
  - Flight Nurse
  - Flight Respiratory Therapist
  - Physician
  - Nurse Practitioner/Physician Assistant
  - Other. Please specify: \_\_\_\_\_
-

Q3 Which of the following best describes your main EMS agency?

- Hospital** - Refers to EMS agencies that are under the direct control of a hospital.
  - Fire Department (Paid and Volunteer)** - An organization from which fire and emergency medical services are provided.
  - Tribal** - An EMS agency operated by a federally recognized Indian or Alaska Native Tribe.
  - Military** - An agency operated by one of the U.S. Armed Forces.
  - Government, Non-Fire Department** - An agency operated directly by a federal, state, county, or local government entity other than the U.S. Armed Forces.
  - Private** - An agency operated under the direct control of a for-profit or not-for-profit organization other than a hospital to include volunteer rescue squads operated independently of a fire department.
  - Air Medical** - An organization which provides air ambulance services.
  - Other.** Please specify: \_\_\_\_\_
-

Q4 Which of the following best describes the primary type of service provided by your main EMS agency? If more than one type of service is provided, pick the service with the greatest number of calls in the past 12 months.

- Air Medical Care** - Transport of patients from scene or hospital by rotor or fixed wing.
- Primarily 911 response** - Immediate response to an incident location, regardless of method of notification (for example, 911, direct dial, walk-in, flagging down).
- Primarily medical transport** - Transport of a patient from one health facility to another.
- Equal mix** of 911 and medical transport.
- Clinical services** - Provision of clinical services in a non-ambulance clinical setting such as emergency department, medical office, or dialysis clinic.
- Mobile Integrated Healthcare & Community Paramedicine**- Provision of clinical services in an out-of-hospital community setting.
- Other.** Please specify: \_\_\_\_\_

End of Block: Introductory Questions- Page 1

---

Start of Block: Introductory Questions- Page 2

Q5 A major trauma patient in your area is taken by ground to:

- Closest Level I or II trauma center
- Closest Level III or IV trauma center
- Closest hospital (non-trauma center)



Q6 What is the average time by ground, in minutes, with good weather and low traffic volumes, to reach your closest **trauma center** of any level?

\_\_\_\_\_

---



Q7 What is the average time by ground, in minutes, with good weather and low traffic volumes, to reach your closest **Level I or II trauma center**?

---

---

Q8 A major trauma patient in your area taken by air:

- 0% of the time
  - 1-25% of the time
  - 26-50% of the time
  - 51-75% of the time
  - 76-99% of the time
  - 100% of the time
- 

Q9 In your region, how often is air transport **unavailable** (e.g. due to weather, aircraft availability)?

- 0% of the time
- 1-25% of the time
- 26-50% of the time
- 51-75% of the time
- 76-99% of the time
- 100% of the time

End of Block: Introductory Questions- Page 2

---

Start of Block: Introductory Questions - Page 3

Q10 When there is new information regarding trauma patient care, how do you typically find out about it before it is integrated into your local protocol? Select all that apply.

- Attending a conference
  - Internet/newspaper/television
  - Academic journals
  - State or regional authorities
  - Local agency
  - Colleagues
  - Social media
  - Other. Please specify: \_\_\_\_\_
- 

Q11 What percentage of the time does your judgement override the FTG?

- 0-20%
  - 21-40%
  - 41-60%
  - 61-80%
  - 81-100%
-

Q12 When your judgement overrides the FTG, do you most often find yourself:

- a. Transporting a patient to a trauma center when they don't necessarily meet criteria based on the current FTG?
- b. Transporting a patient to a non-trauma center when they meet criteria for a trauma center based on FTG?
- c. Equal mix of both a and b.



Q13 What transport time, by ground, in minutes, do you think is reasonable to travel to a Level I or II trauma center as opposed to a closer hospital with a patient meeting Step 1 or 2 triage criteria?

---

End of Block: Introductory Questions - Page 3

---

Start of Block: Introductory Questions - Page 4

Q14 Which Step drives the majority of the decisions to take patients to a trauma center in your area?

- Step 1- Measure vital signs and level of consciousness
  - Step 2- Assess anatomy of injury
  - Step 3- Assess mechanism of injury and evidence of high-energy impact
  - Step 4- Assess special patient or system considerations
-

Q15 Please order the steps in terms of importance to you as a provider in the field determining patient transport destination. (Drag and drop your responses. "1" being most important; "4" being least important)

- \_\_\_\_\_ Assess mechanism of injury and evidence of high-energy impact
  - \_\_\_\_\_ Assess anatomy of injury
  - \_\_\_\_\_ Assess special patient or system considerations
  - \_\_\_\_\_ Measure vital signs and level of consciousness
- 

Q16 Please place the steps in the order you use them as a provider in the field in determining patient transportation destination. (Drag and drop your responses.)

- \_\_\_\_\_ Assess mechanism of injury and evidence of high-energy impact
  - \_\_\_\_\_ Assess anatomy of injury
  - \_\_\_\_\_ Assess special patient or system considerations
  - \_\_\_\_\_ Measure vital signs and level of consciousness
- 

Q17 Do you think this step-wise approach to triage is useful for implementation of trauma triage criteria?

Yes. Why? \_\_\_\_\_

No. Why not? \_\_\_\_\_

---

Q18 Are there any other specific tools you find helpful to assist in consistent implementation of trauma triage criteria?

---

---

---

---

---

End of Block: Introductory Questions - Page 4

---

Start of Block: Step 1: Measure Vital Signs and Level of Consciousness

Step 1 Q1

**Step 1: Measure Vital Signs and Level of Consciousness**

Is **Step 1** of the FTG (physiologic criteria) used in your local protocol?

Yes

No

---

Step 1 Q2 How easy is it to understand **Step 1** of the FTG when used in your local protocol?

Very easy

Easy

Neither easy nor difficult

Difficult. Why? \_\_\_\_\_

Very difficult. Why? \_\_\_\_\_

Not applicable

---

Step 1 Q3 How important is **Step 1** of the FTG to triage?

Extremely important

Very important

Moderately important

Slightly important

Not at all important

---

Step 1 Q4 Please indicate whether you would keep, eliminate, or change each of the **physiologic measures** within Step 1 of the FTG.

	Keep	Eliminate	Change
<input checked="" type="checkbox"/> Glasgow Coma Scale $\leq 13$	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Systolic Blood Pressure	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Respiratory rate 29 breaths per minute ( )	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

*Display This Question:*

*If Please indicate whether you would keep, eliminate, or change each of the physiologic measures wit... = Change*

Step 1 Q4.1 As indicated in the question above, please describe what you would change about the **physiologic measures** within Step 1 of the FTG.

---



---



---



---



---

Step 1 Q5 Are there any additional **physiologic measures** that are used by your EMS agency to support trauma center triage?

- Yes. Please specify: \_\_\_\_\_
- No

Step 1 Q6 Are there any additional **physiologic measures** you believe should be added in the next iteration of the Guidelines?

- Yes. Please specify: \_\_\_\_\_
- No

**End of Block: Step 1: Measure Vital Signs and Level of Consciousness**

---

**Start of Block: Step 2: Assess Anatomy of Injury**

Step 2 Q1 **Step 2: Assess Anatomy of Injury**

Is **Step 2** of the FTG used in your local protocol?

- Yes
- No

-----

Step 2 Q2 How easy is it to understand **Step 2** of the FTG when used in your local protocol?

- Very easy
- Easy
- Neither easy nor difficult
- Difficult. Why? \_\_\_\_\_
- Very difficult. Why? \_\_\_\_\_
- Not applicable
-

Step 2 Q3 How important is **Step 2** of the FTG to triage?

- Extremely important
  - Very important
  - Moderately important
  - Slightly important
  - Not at all important
-

Step 2 Q4 Please indicate whether you would keep, eliminate, or change each ***injury type*** used to identify the major trauma patient?

	Keep	Eliminate	Change
<input checked="" type="checkbox"/> All penetrating injuries to head, neck, torso and extremities proximal to elbow or knee	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Chest wall instability or deformity (e.g. flail chest)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Two or more proximal long-bone fractures	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Crushed, degloved, mangled, or pulseless extremity	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Amputation proximal to wrist or ankle	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Pelvic Fractures	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Open or depressed skull fracture	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> Paralysis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

*Display This Question:*

*If Please indicate whether you would keep, eliminate, or change each injury type used to identify th...  
= Change*

Step 2 Q4.1 As indicated in the question above, please describe what you would change about the ***injury types*** used to identify the major trauma patient.

---

---

---

---

---

---

Step 2 Q5 Are there any additional ***injury types*** that are used by your EMS agency to support trauma center triage?

- Yes. Please specify: \_\_\_\_\_
- No

---

Step 2 Q6 Are there any additional ***injury types*** you believe should be added to improve the next iteration of the Guideline?

- Yes. Please specify: \_\_\_\_\_
- No

End of Block: Step 2: Assess Anatomy of Injury

---

Start of Block: Step 3: Assess Mechanism of Injury and Evidence of High-energy Impact

Step 3 Q1

**Step 3: Assess Mechanism of Injury and Evidence of High-energy Impact**

Is **Step 3** of the FTG used in your local protocol?

- Yes
- No

---

Step 3 Q2 How easy is it to understand **Step 3** of the FTG when used in your local protocol?

- Very easy
  - Easy
  - Neither easy nor difficult
  - Difficult. Why? \_\_\_\_\_
  - Very difficult. Why? \_\_\_\_\_
  - Not applicable
- 

Step 3 Q3 How important is **Step 3** of the FTG to triage?

- Extremely important
  - Very important
  - Moderately important
  - Slightly important
  - Not at all important
-

Step 3 Q4 Please indicate whether you would keep, eliminate, or change each **mechanism of injury** used to identify the major trauma patient?

	Keep	Eliminate	Change
<p><input checked="" type="checkbox"/> <b>Falls</b>- Adults: &gt;20 feet (one story is equal to 10 feet) Children: &gt;10 feet or two or three times the height of the child</p>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<p><input checked="" type="checkbox"/> <b>High-risk auto crash</b>- Intrusion, including roof: &gt;12 inches occupant site; &gt;18 inches any site   Ejection (partial or complete) from automobile   Death in same passenger compartment   Vehicle telemetry data consistent with a high risk of injury</p>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<p><input checked="" type="checkbox"/> <b>Auto vs. pedestrian/ bicyclist</b> thrown, run over, or with significant (&gt; 20 mph) impact</p>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<p><input checked="" type="checkbox"/> <b>Motorcycle crash</b> &gt; 20 mph</p>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Display This Question:

If Please indicate whether you would keep, eliminate, or change each mechanism of injury used to ide... = Change

Step 3 Q4.1 As indicated in the question above, please describe what you would change about the **mechanisms of injury** used to identify the major trauma patient.

---



---

---

---

---

Step 3 Q5 Are there any additional ***mechanisms of injury*** that are used by your EMS agency to support trauma center triage?

- Yes. Please specify: \_\_\_\_\_
- No

Step 3 Q6 Are there any additional ***mechanisms of injury*** you believe should be added to improve the next iteration of the Guideline?

- Yes. Please specify: \_\_\_\_\_
- No

End of Block: Step 3: Assess Mechanism of Injury and Evidence of High-energy Impact

Start of Block: Step 4: Assess Special Patient or System Considerations

Step 4 Q1 **Step 4: Assess Special Patient or System Considerations**

Is **Step 4** of the FTG used in your local protocol?

- Yes
- No

Step 4 Q2 How easy is it to understand **Step 4** of the FTG when used in your local protocol?

- Very easy
  - Easy
  - Neither easy nor difficult
  - Difficult. Why? \_\_\_\_\_
  - Very difficult. Why? \_\_\_\_\_
  - Not applicable
- 

Step 4 Q3 How important is **Step 4** of the FTG to triage?

- Extremely important
  - Very important
  - Moderately important
  - Slightly important
  - Not at all important
-

Step 4 Q4 Please indicate whether you would keep, eliminate, or change each **special patient or system consideration** influencing your identification of a major trauma victim?

	Keep	Eliminate	Change
<input checked="" type="checkbox"/> <b>Older Adults:</b> Risk of injury/death increases after age 55 years   SBP <110 may represent shock after age 65   Low impact mechanisms (e.g. ground level falls) may result in severe injury	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> <b>Children:</b> Should be triaged preferentially to pediatric capable trauma centers	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> <b>Anticoagulants and bleeding disorders:</b> Patients with head injury are at high risk for rapid deterioration	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> <b>Burns:</b> Without other trauma mechanism: triage to burn facility   With trauma mechanism: triage to trauma center	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> <b>Pregnancy &gt;20 weeks</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
<input checked="" type="checkbox"/> <b>EMS provider judgment</b>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Display This Question:

If Please indicate whether you would keep, eliminate, or change each special patient or system consi... = Change

Step 4 Q4.1 As indicated in the question above, please describe what you would change about the **special patient or system considerations** influencing your identification of a major trauma victim.

---

---

---

---

---

Step 4 Q5 Are there any additional **special patient or system considerations** that are used by your EMS agency to support trauma center triage?

- Yes. Please specify: \_\_\_\_\_
- No

Step 4 Q6 Are there any additional **special patient or system considerations** you believe should be added to improve the next iteration of the Guideline?

- Yes. Please specify: \_\_\_\_\_
- No

End of Block: Step 4: Assess Special Patient or System Considerations

---

## Appendix B:

Organizations that were requested to send the end-user feedback tool to their membership.

**Organizations**

American Academy of Orthopedic Surgeons (AAOS)
American Academy of Pediatrics (AAP)
American Ambulance Association (AAA)
American Association for the Surgery of Trauma (AAST)
American Association for the Surgery of Trauma (AAST)
American College of Emergency Physicians (ACEP)
American Trauma Society (ATS)
Association of Air Medical Services (AAMS)
Commission on Accreditation for Prehospital Continuing Education (CAPCE)
Commission on Accreditation of Medical Transport Systems (CAMTS)
Committee on Accreditation of EMS Programs (CoAEMSP)
Eastern Association for the Surgery of Trauma (EAST)
Emergency Medical Services for Children (EMSC)
Emergency Nurses Association (ENA)
International Association of Fire Chiefs (IAFC)
International Association of Fire Fighters (IAFF)
International Trauma Life Support (ITLS)
National Association of Emergency Medical Technicians (NAEMT)
National Association of EMS Educators (NAEMSE)
National Association of EMS Physicians (NAEMSP)
National Association of State 911 Administrators (NASNA)
National Association of State EMS Officials (NASEMSO)
National Registry of Emergency Medical Technicians (NREMT)
National Volunteer Fire Council (NVFC)
Orthopedic Trauma Association (OTA)
Pediatric Trauma Society (PTS)
Prehospital Trauma Life Support (PHTLS)
Society of Trauma Nurses (STN)
Western Trauma Association (WTA)