

Stepped collaborative care for trauma: giant leaps for health equity

Rachel Rivero, Lisa M Kodadek 

To cite: Rivero R, Kodadek LM. Stepped collaborative care for trauma: giant leaps for health equity. *Trauma Surg Acute Care Open* 2024;**9**:e001359. doi:10.1136/tsaco-2024-001359

Mortality rates after injury are low due to advances in trauma care, but survival is not sufficient: trauma leads to more disability worldwide than any other disease.¹ In the year after injury, between 20% and 44.4% of patients screen positive for post-traumatic stress disorder (PTSD) and over 40% of patients are unable to return to work.^{2–4} The goal of a trauma system is to facilitate recovery after injury, including optimal long-term functional outcomes and reintegration into society. However, patients recovering from injury face significant barriers to receiving optimal postdischarge healthcare. Only 28% of level 1 and 2 trauma centers in the USA routinely screen for PTSD⁵ and less than 10% of patients receive treatment for PTSD after injury.³ Patients belonging to minority populations are disproportionately affected by care transition disruption and experience adverse healthcare disparities after trauma.⁶

Dr Abu⁷ and colleagues offer an innovative, stepped collaborative care intervention which has differentially reduced PTSD symptoms at 6 months after injury in self-reported racial and ethnic minority populations. Their work demonstrates the development of a successful model for achieving more equitable health outcomes after injury. These findings result from secondary analysis of data from a stepped wedge, cluster randomized trial conducted at 25 trauma centers. Injured patients at risk for PTSD were randomized to enhanced usual care or the intervention, which included proactive care management, cognitive-behavioral therapy, and psychopharmacology for PTSD.

The authors demonstrate how multidisciplinary care can be used to mitigate health disparities in trauma by incorporating shared decision-making and individual treatment preferences. The collaborative approach highlights the importance of engagement from all healthcare providers, including nurses, social workers, and physicians. Effective care strategies must incorporate patient-centered care and tailored elements to directly address baseline healthcare inequities and social determinants of health. Furthermore, care must account for patients' individual past experiences, their own interpretation of trauma, and culturally specific ways of healing.

All patients included in this study had very high rates of baseline PTSD and prior trauma. Although we often think of trauma as an acute disease, the reality is that it may be recurrent and chronic, particularly when associated behavioral health risk factors are not addressed. Among non-white/Hispanic patients, 14.3% had baseline PTSD and almost 40% had suffered five or more prior injuries; they also reported significantly lower frequencies of preinjury mental health service utilization compared with white/non-Hispanic patients. For

some, the collaborative care intervention may have represented the first opportunity to receive appropriate healthcare for PTSD associated with prior injury.

Dr Abu and colleagues should be commended for taking important steps to care for adverse sequelae associated with traumatic injury. For underserved populations, these first steps represent giant leaps for health equity.

Contributors RR and LMK drafted the commentary. LMK and RR provided critical revisions and approved the final commentary.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Lisa M Kodadek <http://orcid.org/0000-0001-6433-9159>

REFERENCES

- Vos T, Lim SS, Abbafati C, Abbas KM, Abbasi M, Abbasifard M, Abbasi-Kangevari M, Abbastabar H, Abd-Allah F, Abdelalim A, *et al*. Diseases and injuries collaborators. global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the global burden of disease study 2019. *The Lancet* 2020;396:1204–22.
- Haider AH, Herrera-Escobar JP, Al Rafai SS, Harlow AF, Apoj M, Nehra D, Kasotakis G, Brasel K, Kaafarani HMA, Velmahos G, *et al*. Factors associated with long-term outcomes after injury results of the functional outcomes and recovery after trauma emergencies (FORTE) multicenter cohort study. *Ann Surg* 2020;271:1165–73.
- Bell TM, Vektor AN, Zarzaur BL. Prevalence and treatment of depression and Posttraumatic stress disorder among trauma patients with non-neurologic injuries. *J Trauma Acute Care Surg* 2018;85:999–1006.
- deRoos-Cassini TA, Hunt JC, Geier TJ, Warren AM, Ruggiero KJ, Scott K, George J, Halling M, Jurkovich G, Fakhry SM, *et al*. Screening and treating hospitalized trauma survivors for PTSD and depression. *J Trauma Acute Care Surg* 2019;87:440–50.
- Bulger EM, Johnson P, Parker L, Moloney KE, Roberts MK, Vaziri N, Seo S, Nehra D, Thomas B, Zatzick D. Nationwide survey of trauma center screening and intervention practices for Posttraumatic stress disorder, firearm violence. *J Am Coll Surg* 2022;234:274–87.
- Chun Fat S, Herrera-Escobar JP, Seshadri AJ, Al Rafai SS, Hashmi ZG, de Jager E, Velmahos C, Kasotakis G, Velmahos G, Salim A, *et al*. Racial disparities in post-discharge healthcare utilization after trauma. *Am J Surg* 2019;218:842–6.
- Abu K, Bedard-Gilligan M, Moodliar R, Bulger EM, Hernandez A, Knutzen T, Shoyer J, Birk N, Conde C, Engstrom A, *et al*. Can stepped collaborative care interventions improve post-traumatic stress disorder symptoms for racial and ethnic minority injury survivors? *Trauma Surg Acute Care Open* 2024;9:e001232.



► <http://dx.doi.org/10.1136/tsaco-2023-001232>

© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Yale School of Medicine, New Haven, Connecticut, USA

Correspondence to

Dr Lisa M Kodadek; lisa.kodadek@yale.edu