

# Early identification of respiratory decompensation among older adults with rib fractures: a sound solution for fragile ribs

W Preston Hewgley, Adam Lucy, Rondi Gelbard 

**To cite:** Hewgley WP, Lucy A, Gelbard R. Early identification of respiratory decompensation among older adults with rib fractures: a sound solution for fragile ribs. *Trauma Surg Acute Care Open* 2024;9:e001632. doi:10.1136/tsaco-2024-001632  
Accepted 9 October 2024

Failure to rescue in the setting of respiratory decline contributes significantly to unplanned intensive care unit admissions (UIAs). In this context, Davis *et al*<sup>1</sup> evaluate a novel continuous monitor of pulmonary function (Integrated Pulmonary Index, IPI) to identify early signs of respiratory decompensation among older adults with rib fractures. In this thought-provoking study, the authors found that UIAs were significantly lower among patients monitored with IPI. In our opinion, a critical examination of this study is warranted to contextualize their findings and help guide future research on this important topic. Here we highlight a few areas of importance.

Predicting which patients are likely to clinically deteriorate is important for mitigating poor outcomes. Whereas the IPI was intended to identify early signs of respiratory decompensation, it is unclear which patients are most likely to benefit from this tool. The retrospectively selected control group appears to have a very different baseline risk profile than the intervention cohort, which was older and had higher rates of smoking, heart failure or stroke. It is unclear if the authors controlled for these differences in their analysis. Moreover, providers were not required to implement IPI monitoring on all eligible patients, perhaps introducing selection bias into their sample population.

In addition, UIAs increase overall length of stay and resource utilization and are an important quality metric in trauma care. Patients with rib fractures are especially vulnerable to unanticipated transfer, partially due to difficulty in initial triaging. However, the IPI tool was not intended to help risk stratify patients or guide initial triage disposition. Several risk-based triage tools have been studied in this capacity (ie, Rib Injury Guidelines Score, PIC Score, Forced Vital Capacity), and as such, may be better suited for informing admission triage and to downgrade decisions.<sup>2-4</sup> Although the IPI tool appears to be a useful adjunct for alerting providers

when a patient unexpectedly deviates from the predicted clinical trajectory, it seems that appropriate triage at admission and effective protocols to optimize care are also fundamental to improving outcomes in this vulnerable patient population.

**Contributors** RG: conception, writing, and editing; WPH: conception, writing, and editing; AL: conception, writing, and editing.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** Not applicable.

**Provenance and peer review** Commissioned; internally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Rondi Gelbard <http://orcid.org/0000-0003-3613-4943>

#### REFERENCES

- 1 Davis N, Lindbloom P, Hromatka K, Gipson J, West MA. Use of an Integrated Pulmonary Index pathway decreased unplanned ICU admissions in elderly patients with rib fractures. *Trauma Surg Acute Care Open* 2024;9:e001523.
- 2 Nelson A, Reina R, Northcutt A, *et al*. Prospective validation of the Rib Injury Guidelines for traumatic rib fractures. *J Trauma Acute Care Surg* 2022;92:967–73.
- 3 Witt CE, Bulger EM. Comprehensive approach to the management of the patient with multiple rib fractures: a review and introduction of a bundled rib fracture management protocol. *Trauma Surg Acute Care Open* 2017;2:e000064.
- 4 Billings JD, Khan AD, Clement LP, *et al*. A clinical practice guideline using percentage of predicted forced vital capacity improves resource allocation for rib fracture patients. *J Trauma Acute Care Surg* 2021;90:769–75.



► <http://dx.doi.org/10.1136/tsaco-2024-001523>

© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Surgery, The University of Alabama at Birmingham Heersink School of Medicine, Birmingham, Alabama, USA

#### Correspondence to

Dr Rondi Gelbard; [rgelbard@uabmc.edu](mailto:rgelbard@uabmc.edu)