Supplemental Table 1. Differences Between Border-Fence Falls and Domestic Falls by Injury

Characteristics, n (%)	Border, N=64 (52%)	Domestic, N=60 (48%)	P-value
Head	2 (3%)	15 (25%)	0.004
Intracranial hemorrhage	2 (3%)	13 (22%)	0.002
Subdural hemorrhage	1 (2%)	9 (15%)	0.007
Subarachnoid hemorrhage	1 (2%)	10 (17%)	0.003
Intraparenchymal hemorrhage	1 (2%)	1 (2%)	>0.99
Soft tissue injury	0 (0%)	6 (10%)	0.01
Face	1 (2%)	7 (12%)	0.03
Maxillary fracture	0 (0%)	1 (2%)	0.48
Mandibular fracture	1 (2%)	0 (0%)	>0.99
Zygomatic fracture	0 (0%)	3 (5%)	0.11
Nasal bone fracture	0 (0%)	3 (5%)	0.11
Orbital fracture	0 (0%)	4 (7%)	0.05
Contusion	0 (0%)	3 (5%)	0.11
Eye injury	0 (0%)	1 (2%)	0.48
Chest	3 (5%)	16 (27%)	0.007
Rib fracture	2 (3%)	13 (22%)	0.002
Lung contusion	1 (2%)	9 (15%)	0.007
Pneumothorax	2 (3%)	7 (12%)	0.09
Hemothorax	0 (0%)	4 (7%)	0.05
Clavicular fracture	1 (2%)	3 (5%)	0.35
Scapular fracture	0 (0%)	3 (5%)	0.11
Abdomen	0 (0%)	4 (7%)	0.05
Kidney laceration	0 (0%)	2 (3%)	0.23
Splenic laceration	0 (0%)	1 (2%)	0.48
Liver laceration	0 (0%)	2 (3%)	0.23
Spine	23 (36%)	21 (35%)	0.91
Cervical fracture	0 (0%)	4 (7%)	0.05
Lumbar fracture	18 (28%)	17 (28%)	0.98
Thoracic fracture	9 (14%)	8 (13%)	0.90
Pelvis	4 (6%)	6 (10%)	0.52
Ilium fracture	1 (2%)	3 (5%)	0.35
Rami fracture	4 (6%)	5 (8%)	0.74
Sacral fracture	4 (6%)	2 (3%)	0.68
Acetabular fracture	1 (2%)	4 (7%)	0.20
Extremities	47 (73%)	25 (42%)	0.003
Ankle fracture	12 (19%)	6 (10%)	0.17
Foot fracture	2 (3%)	2 (3%)	>0.99
Fibular fracture	22 (34%)	7 (12%)	0.003

Hand avulsion	2 (3%)	2 (3%)	>0.99	
Calcaneus fracture	14 (22%)	6 (10%)	0.07	
Tibial fracture	24 (38%)	9 (15%)	0.005	
Humeral fracture	1 (2%)	4 (7%)	0.20	
Ulnar fracture	0 (0%)	5 (8%)	0.02	
Radial fracture	4 (6%)	9 (15%)	0.11	
Femur fracture	1 (2%)	3 (5%)	0.35	
Elbow dislocation	1 (2%)	1 (2%)	>0.99	
Shoulder dislocation	0 (0%)	1 (2%)	0.48	
Knee contusion	1 (2%)	0 (0%)	>0.99	